

Erratum to: Antihypertensive Efficacy of Triple Combination Perindopril/Indapamide Plus Amlodipine in High-Risk Hypertensives: Results of the PIANIST Study (Perindopril-Indapamide plus Amlodipine in high risk hypertensive patients)

Kálmán Tóth · on behalf of PIANIST Investigators

Published online: 11 April 2014
© Springer International Publishing Switzerland 2014

**Erratum to: Am J Cardiovasc Drugs (2014) 14:137–145
DOI 10.1007/s40256-014-0067-2**

Page 137, abstract, paragraph 3, lines 3 and 4 which previously read as:

that was not properly controlled despite antihypertensive therapy, and for whom study treatment

should read as:

that was not properly controlled, and for whom study treatment

Page 138, column 2, paragraph 2, line 4 which previously read:

treatment and were at high or very high cardiovascular risk

should read:

treatment and/or were at high or very high cardiovascular risk

Page 141, column 1, paragraph 1, line 1 which previously read:

treatments (Fig. 2), 81 and 91 % of patients previously should read:

treatments (Fig. 2), 91 and 81 % of patients previously

The online version of the original article can be found under doi:[10.1007/s40256-014-0067-2](https://doi.org/10.1007/s40256-014-0067-2).

K. Tóth (✉)
1st Department of Medicine, University of Pécs, Medical School, Ifjúság St. 13, Pécs 7624, Hungary
e-mail: Kalman.Toth@aok.pte.hu; toth.kalman@pte.hu