

S0031-9422(96)00132-X

A 1,5-CIS-GUAIANOLIDE GLUCOSIDE FROM FLOWERS OF ARNICA MOLLIS

IRMGARD MERFORT,*† GÜNTER WILLUHN,* DETLEF WENDISCH‡ and DANIEL GONDOL‡

*Institut für Pharmazeutische Biologie, Heinrich-Heine-Universität Düsseldorf, Geb. 26.23, Universitätsstr. 1, D-40225 Düsseldorf, Germany: ‡Abteilung ZF-D Strukturforschung, Bayer AG Leverkusen, D-51368 Leverkusen, Germany

(Received 11 December 1995)

Key Word Index—*Arnica mollis*; Asteraceae; sesquiterpene lactone glucoside; guaianolide; pinoresinol glucoside; benzyl glucoside.

Abstract—The novel sesquiterpene lactone glucoside $2\alpha - O - \beta$ -D-glucopyranosyl- 4α -hydroxy- 1α , 5α , 11α -H-guai-10(14)-en-12, 8α -olide (= 2α -hydroxy- 11α ,13-dihydro-1-epi-inuviscolide- 2α -O- β -D-glucopyranoside) was iso-lated from flowerheads of *Arnica mollis*. Its structural elucidation was carried out by NMR spectroscopy (¹H and ¹³C NMR, ¹H–¹H COSY, HMBC, HMQC and ¹H–¹H NOESY) and mass spectrometry (DCI). Additionally, the lignan (+)-pinoresinol-4'- β -D-glucopyranoside and benzyl- β -glucoside were isolated and identified.

INTRODUCTION

Previously we reported on the isolation and identification of two xanthanolide glucosides from flowerheads of Arnica anplexicaulis and A. mollis, which have been the first sesquiterpene lactone glycosides detected within the genus Arnica [1]. In continuation of our chemical study of this genus [2] we have now isolated a guainolide glucoside from a methanolic extract of the flowers from A. mollis, which, to the best of our knowledge, is described for the first time. The aglycone is also a new compound. Additionally, the lignan, (+)-pinoresinol-4'- β -D-glucopyranoside, and benzyl- β glucoside were found in the plant.

RESULTS AND DISCUSSION

The DCI (NH₃) mass spectrum of the isolated sesquiterpene lactone glycoside displayed a quasimolecular ion at m/z 446 [M + NH₄]⁺ and fragment ions at m/z 428 [M - H₂O + NH₄]⁺, 284 [M - 162 + NH₄]⁺ and 266 [M - H₂O - 162 + NH₄]⁺, due to the elimination of H₂O and/or a hexose, from which a molecular mass of 426 and the molecular formula $C_{21}O_9H_{32}$ could be deduced. Acid hydrolysis on a TLC plate [3] gave glucose as the sugar moiety.

Structural elucidation was achieved by ¹H and ¹³C NMR spectroscopy, including ¹H–¹H COSY, HMQC and HMBC, which allowed the full assignment of all carbon and proton resonances. Additionally, the stereo-chemistry was determined by a ¹H–¹H NOESY spec-

trum. The ¹³C and ¹H NMR spectra (Tables 1 and 2) show the typical signals for a β -D-glucopyranose with the doublet for H-1' at δ 4.27 and the resonance for C-1' at δ 102.5, being linked to a secondary hydroxyl group at the aglycone [4, 5]. The remaining 'H NMR signals of the aglycone are in part similar to those of the 1,5-cis-guainolide, 11 β ,13-dihydro-1-epi-inuviscolide, from Dittrichia graveolens [6]. Marked difference exist only in the chemical shifts and coupling patterns of the protons H-2, H-3 and H-5, which are explainable by the assumption that the additional Oglucoside moiety is located at C-2. Furthermore, the signals for H-7, H-8 and H-11 are shifted downfield (0.52, 0.28 and 0.39 ppm, respectively) compared to those of the above mentioned 11β , 13-dihydro-derivative, suggesting an 11β -oriented methyl group for which such downfield shifts are characteristic [1, 7, 8]. In the ¹³C NMR spectrum (Table 2) the signal for C-12 is located at δ 182.3, which is typical for a saturated lactone ring, indicating the absence of an exo-methylene group at C-11 [9]. The resonance of C-6 at δ 23.5 and C-13 at δ 9.9, which show a correlation with the signal for H-11 (at δ 2.66) in the HMBC spectrum, as well as that for C-13, correspond with those for C-6, C-11 and C-13 of the guaianolide 11α , 13-dihydro-2-Oacetyl-florilenalin (C-6: δ 22.9; C-11: δ 41.0, C-13: δ 9.9 [8]). From the HMQC spectrum the occurrence of two further quaternary carbons with δ 144.9 and 86.0 are deducible. Due to their chemical shifts and literature data [8] they have to be assigned to C-10 and C-4, respectively. Moreover, the HMBC spectrum exhibited long range couplings between the methylene protons H-14 and C-1 as well as C-9. The attachment of glucose to the hydroxyl at C-2 is confirmed by the long range coupling between H-1' and C-2.

[†]Address for correspondence: Institut für Pharmazeutische Biologie, Albert-Ludwigs-Universität Freiburg, Schänzlestr. 1, 79104 Freiburg, Germany.

Table 1. ¹H NMR data for compound 1 (500 MHz, CD₃OD, TMS as internal standard)*

Proton	δ (ppm)		J (Hz)	Correlated with H	NOE with H [†]
1	2.99	dd	11.8, 6.6	2,5	$5, 7, 9\alpha, 1'$
2	4.63‡	ddd	11.8, 9.3, 7.1	$1, 3\alpha, 3\beta$	3 <i>B</i>
3α	1.99	dd	9.3, 12.1	2, 3 <i>B</i>	3 ^β , 5
3 <i>B</i>	2.16	dd	7.1, 12.1	$2, 3\alpha$	2β , 15-H ₁ , 3α
5	2.43‡	m	6.6, 4.8, 13.6	$1, 6\alpha, 6\beta$	1, 6α, 7
6α	1.78	dd(br)	4.8, 13.6	5, 6 <i>β</i>	5, 6 <i>β</i> , 7
6 <i>β</i>	1.07	ddd	13.6, 13.6, 10.6	$5, 6\alpha, 7$	6 <i>α</i> , 8
7	2.19-2.27	т	10.6, 10.8	$6\beta, 8$	$1, 5, 6\alpha, 11$
8	4.07	ddd	10.8, 10.8, 4.9	$7, 9\alpha, 9\beta$	6 <i>β</i> , 9 <i>β</i> , 13-Η,
9α	2.19-2.27	т	10.8, 11.5	8, 9 <i>β</i>	9β
9β	3.06	dd	4.9, 11.5	$8,9\alpha$	8, 9 <i>α</i> , 14b
11	2.66	dq§	7.7, 7.7	7,13	7, 13-H,
13	1.15	d	7.7	11	8, 11
14a	5.28	5			
14b	5.18	S			
15	1.08	S			
1'	4.27	d	7.7	2'	
2'	3.11-3.15	m	8.8	3'	
3'	3.28	t	8.8, 8.8	2', 4'	
4'	3.23	t	8.8	3'	
5'	3.11-3.15	т	2.3, 6.0	6a', 6b'	
6a'	3.85	dd	2.3, 11.9	5, 6b'	
6b'	3.64	dd	6.0, 11.9	5, 6a'	

*Assignments are based on ${}^{1}H-{}^{1}H$ COSY.

†From ¹H-¹H NOESY.

‡Not first-order signal.

§Pseudo-quintet.

The α -configuration of H-1 and thus the evidence for a $1_{\alpha}, 5_{\alpha}$ -cis-guaianolide is given from ${}^{3}J_{1\alpha,5\alpha} = 6.6$ Hz [7, 10], whereas in the case of a $1\beta, 5\alpha$ -transguaianolide a value of 10 Hz or greater should be anticipated [11, 12]. This stereochemistry is confirmed by strong nOe effects between H-1 and H-5.

The configuration at C-2 (2 β -H) follows from ${}^{3}J_{1\alpha,2\beta} = 11.8$ Hz and the lacking of a nOe correlation

Table 2. ¹³C NMR data for compound 1 (125 MHz, CD₃OD, TMS as int. standard)*

A	glycone	Sugar		
Carbon	δ (ppm)	Carbon	δ (ppm)	
1	53.5	1'	102.5	
2	78.5	2'	75.3	
3	ca 48.5†	3'	78.7	
4	86.0	4'	72.0	
5	54.6	5'	78.5	
6	23.5	6'	62.8	
7	47.0			
8	78.5			
9	44.9			
10	144.9			
11	41.8			
12	182.3			
13	9.9			
14	116.6			
15	24.4			

*Assignments are based on HMBC and HMQC.

†Hidden under solvent signals.

between H-1 and H-2. Clear nOe correlations are observed between H-7 and H-11 as well as H-8 and 13-H₃, confirming the 11 β -position of the methyl group at C-13. The absence of nOe correlation between H-7 and H-8 is in accordance with the $7\beta(12),8\alpha$ *trans*-fusion of the lactone ring (${}^{3}J_{7,8} = 10.8$ Hz). The methyl group at C-4 is β -orientated, because no nOe correlation exists with H-5. Hence, the structure of the isolated sesquiterpene lactone is $2\alpha - O - \beta - D$ -glucopyranosyl- 4α -hydroxy- $1\alpha,5\alpha,11\alpha$ -*H*-guai-10(14)-en- $12,8\alpha$ -olide (= 2α -hydroxy- $11\alpha,13$ -dihydro-1-epiinuvisculide- 2α -O- β -D-glucopyranoside (1).

Additionally, benzyl- β -glucoside (2) [13, 14] and the lignan glycoside (+)-pinoresinol-4'- β -D-glucopyranoside (3) [15, 16] were isolated and the structures confirmed by their DCI mass spectral, ¹H NMR and



(lignan) ¹³C NMR and optical rotation data. Within the genus *Arnica* non-glycosylated (+)-pinoresinol has already been found in flowers of *A. angustifolia* [17] and roots of *A. chamissonis* ssp. *foliosa* [13] and benzyl- β -glucoside in the roots of the latter species [13].

EXPERIMENTAL

Plant material. See ref. [7]. A voucher specimen (No. 8.2) is deposited at the herbarium of the Institute for Pharmaceutical Biology, University of Düsseldorf.

Instrumentation. NMR: Bruker AMX 500, 500 MHz (¹H NMR) and 125 MHz (¹³C NMR) in CD₃OD, TMS as int. standard; MS: DCl, Finnigan INCOS 50, NH₃ as reactant gas (emitter heating rate 10 mA s⁻¹, calibration with FC43); TCL: silica gel 60_{F254} , EtOAc-HOAc-HCO₂H-H₂O (100:11:11:27), detection: anisalde-hyde-H₂SO₄; sugar: silica gel 60_{F254} , EtOAc-MeOH-HOAc-H,O (12:3:3:2), detection: thymol-H₂SO₄.

Extraction and purification of compounds. Dried and powdered flowerheads (2.02 kg) were exhaustively extracted with CHCl₃ (162 g) and then with MeOH (395.6 g). A 30-g portion of the MeOH extract was sepd by CC on Sephadex LH-20 with MeOH vielding a fr. with 1 and 2 (200 mg) and a fr. with 3 (700 mg). Sepn of the first fr. by CC on Sephadex LH-20 with cyclohexane-CH₂Cl₂-MeOH (4:7:1) followed by (4:7:2) gave a fr. with 2 (26 mg) from which 2 was isolated by CC on silica gel with EtOAc-n-hexane (10:19) and a fr. with 1 (18 mg), which was chromatographed on silica gel with mixts of EtOAc-n-hexane of increasing polarity, followed by EtOAc and finally by EtOAc-Me₂CO (4:1). CC of the fr. containing 3 on Sephadex LH-20 with cyclohexane-CH₂Cl₂-MeOH (4:7:1) and subsequently (4:7:2) yielded 3. Isolated amounts: 1, 5 mg; 2, 4 mg; 3, 2.5 mg.

 $2\alpha - O - \beta - D - Glucopyranosyl - 4\alpha - hydroxy - 1\alpha, 5\alpha, 11\alpha$ - guai - 10(14) - en - 12, 8\alpha - olide (1). Amorphous: R_j : 0.32; anisaldehyde - H₂SO₄: violet turning through blue to green; MS (DCI) m/z (rel. int.): 446 [M + NH₄]⁺ (100), 428 [M - H₂O + NH₄]⁺ (3), 284 [M - 162 + NH₄]⁺, (11), 266 [M - H₂O - 162 + NH₄]⁺ (47), 180 [hexose - H₂O + NH₄]⁺ (24), 162 [hexose - 2H₂O + NH₄]⁺ (9) ⁻¹H NMR: Table 1; ⁻¹³C NMR: Table 2.

Benzyl-β-glucoside (2). Amorphous; R_f : 0.60; anisaldehyde-H₂SO₄; green; MS (DCI) m/z (rel. int.): 288 [M + NH₄]⁺ (100), 270 [M]' (5), 180 [hexose -H₂O + NH₄]⁺ (14), 162 [hexose - 2H₂O + NH₄]⁺ (5). ¹H NMR: refs [13, 14].

Compound **3.** Amorphous; R_f : 0.38; anisaldehyde-H₂SO₄: pink to violet; MS (DCI) m/z (rel. int.): 538 [M + NH₄]⁺ (37), 376 [M - hexose + NH₄]⁺ (22), 180 [hexose - H₂O + NH₄]⁺ (100), 162 [hexose -2H₂O + NH₄]⁺ (35). ¹H and ¹³C NMR: refs [15, 16]. [α]²⁰_D ca + 6 (EtOH). Acknowledgements—We are grateful to Dr U. Matthiesen (Institut für Klinische Chemie und Laboratoriumsdiagnostik, Heinrich-Heine-Universität Düsseldorf) for the DCI mass spectra and Mrs E. Müller for experimental assistance.

REFERENCES

- Paßreiter, C. M., Merfort, I., Bestendonk, C., Willuhn, G. and Steigel, A. (1996) *Planta Med.* 62, 39.
- Willuhn, G., Merfort, I., Paßreiter, C. M. and Schmidt, T. J. (1995) in *Advances in Compositae Systematics* (Hind, D. J. N., Pope, G. V. and Jeffrey, C., eds), Vol. 1, p. 167. The Royal Botanic Gardens, Kew.
- 3. Merfort, I. and Wendisch, D. (1987) *Planta Med.* 53, 434.
- 4. Wehrli, F. W. and Nishida, T. (1979) Prog. Chem. Org. Nat. Prod. 1.
- Kasai, R., Suzuo, M., Asakawa, J. and Tanaka, O. (1977) Tetrahedron Letters 2, 175.
- Rustaiyan, A., Jakupovic, J., Chau-Thi, T. V., Bohlmann, F. and Sadjadi, A. (1987) *Phytochemistry* 26, 2603.
- Marcinek-Hüpen-Bestendonk, C., Willuhn, G., Steigel, A., Wendisch, D., Middelhauve, B., Wiebcke, M. and Mootz, D. (1990) *Planta Med.* 56, 104.
- Schmidt, T. J., Willuhn, G., Steigel, A. and Wendisch, D. (1995) *Planta Med.* 61, 544.
- Youssef, D. and Frahm, A. W. (1994) *Planta Med.* 60, 572.
- Bohlmann, F. and Zdero, C. (1978) *Phytochemistry* 17, 2032.
- Zdero, C., Bohlmann, F., King, R. M. and Robinson, H. (1986) *Planta Med.* 52, 22.
- Zdero, C., Bohlmann, F., King, R. M. and Robinson, H. (1987) *Phytochemistry* 26, 1207.
- Stausberg, S. (1994) Thesis, University of Düsseldorf, Germany.
- Miyase, T., Ueno, A., Takizawa, N., Kobayashi, H. and Karasawa, H. (1987) *Chem. Pharm. Bull.* 35, 1109.
- van de Velde, V., Lavie, D., Gottlieb, H. E., Perold, G. W. and Scheinmann, F. (1984) J. Chem. Soc., Perkin Trans. I 1159.
- Chiba, M., Okabe, K., Hisada, S., Shima, K., Takemoto, T. and Nishibe, S. (1979) *Chem. Pharm. Bull.* 27, 2868.
- Schmidt, T. J., Willuhn, G., Wendisch, D. and Steigel, A. (1993) *Planta Med.* **59** (Suppl. 1), A 598.