

MOLECULAR AND DEVELOPMENTAL NEUROSCIENCE

Sodium/bicarbonate cotransporter NBCn1/slc4a7 increases cytotoxicity in magnesium depletion in primary cultures of hippocampal neurons

Deborah S. Cooper,¹ Han Soo Yang,¹ Peijian He,² Eunjin Kim,¹ Ira Rajbhandari,¹ Chris C. Yun² and Inyeong Choi¹

¹Department of Physiology, Emory University School of Medicine, 615 Michael Street, Atlanta, GA 30322, USA

²Digestive Disease Division, Department of Medicine, Emory University School of Medicine, 615 Michael Street, Atlanta, GA 30322, USA

Keywords: acid, ischemia, Na/HCO₃ transporter, pH, SLC4A7

Abstract

Growing evidence suggests that pharmacological inhibition of Na/H exchange and Na/HCO₃ transport provides protection against damage or injury in cardiac ischemia. In this study, we examined the contribution of the sodium/bicarbonate cotransporter NBCn1 (slc4a7) to cytotoxicity in cultured hippocampal neurons of rats. In neurons exposed to extracellular pH (pH_o) ranging from 6.2 to 8.3, NBCn1 protein expression increased by fivefold at pH < 6.5 compared to the expression at pH_o 7.4. At pH_o 6.5, the intracellular pH of neurons was ~1 unit lower than that at pH 7.4. Immunocytochemistry showed a marked increase in NBCn1 immunofluorescence in plasma membranes and cytosol of the soma as well as in dendrites, at pH_o 6.5. NBCn1 expression also increased by 40% in a prolonged Mg²⁺-free incubation at normal pH_o. Knockdown of NBCn1 in neurons had negligible effect on cell viability. The effect of NBCn1 knockdown on cytotoxicity was then determined by exposing neurons to 0.5 mM glutamate for 10 min and measuring lactate dehydrogenase (LDH) release from neurons. Compared to normal incubation (pH_o 7.2 for 6 h) after glutamate exposure, acidic incubation (pH_o 6.3 for 6 h) reduced cytotoxicity by 75% for control neurons and 78% for NBCn1-knockdown neurons. Thus, both controls and knockdown neurons showed acidic protection from cytotoxicity. However, in Mg²⁺-free incubation after glutamate exposure, NBCn1 knockdown progressively attenuated cytotoxicity. This attenuation was unaffected by acidic preincubation before glutamate exposure. We conclude that NBCn1 has a dynamic upregulation in low pH_o and Mg²⁺ depletion. NBCn1 is not required for acidic protection, but increases cytotoxicity in Mg²⁺-free conditions.

Introduction

The acid–base transporters involving Na/H exchange and Na/HCO₃ transport are known to affect damage and injury in non-neuronal tissues. In the heart, inhibition or knockout of Na/H exchange protects myocytes against ischemia (Karmazyn, 1999; Wang *et al.*, 2003). Inhibition of Na/HCO₃ transport reduces myocardial ischemia–reperfusion injury (Khandoudi *et al.*, 2001). In astrocytes, inhibition of Na/HCO₃ transport reduces cell vulnerability to acid injury (Giffard *et al.*, 2000). Thus, inhibition of these proteins reduces cell damage in cardiac tissues and astrocytes.

Na/H exchangers and Na/HCO₃ transporters are also present in neurons (Chesler, 2003). However, it is currently unclear whether these proteins are involved in neuronal damage. The primary function of Na/H exchangers and Na/HCO₃ transporters is to extrude acid equivalents from the cytosol and recover intracellular pH (pH_i) from acidification (Romero *et al.*, 2004; Bobulescu *et al.*, 2005). Intracel-

lular acidification, together with extracellular acidification, occurs in global and focal ischemia (Lipton, 1999). Severe acidification is deleterious to neurons (Nedergaard *et al.*, 1991). However, moderate to mild acidification has been shown to protect neurons from damage *in vitro* (Lipton, 1999). Thus, neurons can either undergo or be protected from ischemia-induced damage depending upon the magnitude of acidification.

Ionic changes other than acidification occur during ischemia and, among those changes, an aberrant Mg²⁺ homeostasis is of particular interest. Mg²⁺ serves as a cofactor for ATP, a modulator of ion channels and receptors, and a cofactor for enzymatic reactions. Low Mg²⁺ concentrations in the cerebrospinal fluid and serum are found during the early hours after ischemic stroke (McKee *et al.*, 2005). The magnitude of Mg²⁺ decrease reflects the severity of neuronal damage, and there is a close relationship between low Mg²⁺ levels and viability of neurons under pathological states.

Acid extrusion involving Na/HCO₃ cotransport in neurons is known to be governed by the Na⁺-driven Cl/HCO₃ exchanger, which moves Na⁺ and HCO₃[−] into the cytosol in exchange for cytosolic Cl[−] (Schwiening & Boron, 1994; Baxter & Church, 1996). Recent

Correspondence: Dr Inyeong Choi, as above.

E-mail: ichoi@emory.edu

Received 21 February 2008, revised 2 December 2008, accepted 5 December 2008

molecular and cellular studies suggests that some neurons also express the Na/HCO₃ cotransporter, which moves HCO₃⁻ into the cytosol together with Na⁺ (Cooper *et al.*, 2005; Rickmann *et al.*, 2007; Chen *et al.*, 2008). In prenatal and natal hippocampal neurons, the electroneutral Na/HCO₃ cotransporter (NBCn1), encoded by solute carrier family transporter 4A (SLC4A7), is predominantly localized in the dendrites and soma (Cooper *et al.*, 2005) whereas the Na/HCO₃ cotransporter NBCn2 (SLC4A10) and the Na⁺-driven Cl/HCO₃ exchanger NDCBE (SLC4A8) are mostly localized in the soma (Grichchenko *et al.*, 2001; Giffard *et al.*, 2003; Boedtker *et al.*, 2008). In the present study, we examined NBCn1 expression in primary cultures of embryonic rat hippocampal neurons in acidic pH and Mg²⁺-free incubation. We also examined the contribution of NBCn1 to cytotoxicity of neurons under this aberrant ion homeostasis. We focused our research on NBCn1 in this study, as the transporter is extremely sensitive to cellular and systemic pH changes (Kwon *et al.*, 2002; Jakobsen *et al.*, 2004).

Materials and methods

All experiments were conducted under the NIH guidelines for research on animals, and experimental protocols were approved by the Institutional Animal Care and Use Committee at Emory University.

Primary cultures of embryonic hippocampal neurons

Primary cultures of hippocampal neurons were prepared from fetal E19 Sprague Dawley rats (300 g; Harlan Laboratories, Indianapolis, IN, USA). Pregnant rats were fully anesthetized with isoflurane and quickly decapitated, and fetuses were surgically removed. Fetal brains were removed after decapitation and placed in sterile ice-cold dissecting solution (Eagle's balanced salt solution; Invitrogen, Carlsbad, CA, USA) with 10 mM HEPES and 1 mM sodium pyruvate. Hippocampi were removed under a dissecting microscope and then subjected to enzymatic digestion with papain (Worthington Biochemical Corporation; Lakewood, NJ, USA) as well as mechanical digestion. Neurons were plated in Dulbecco's modification of Eagle's medium (DMEM) for acidification experiments, or Neurobasal medium (Invitrogen) for knockdown experiments and immunocytochemistry. Neurons were incubated in a 5% CO₂ humidified chamber at 37°C for 10–14 days. For acidic (or alkaline) incubation, neurons were incubated for 1 h in media of a series of pH ranging from 6.2 to 8.3. The media contained DMEM (no HCO₃⁻) supplemented with 5% fetal bovine serum, 25 U/mL of combined penicillin and streptomycin, 0.6% dextrose, 0.1% MITO (BD Biosciences, San Jose, CA) serum extender and 5 μM cytosine arabinoside. The pH was adjusted by varying HCO₃⁻ levels according to the Henderson–Hasselbalch equation with the solubility coefficient of 0.03 mmol CO₂/mmHg and P_{CO2} bubbled at 35.65 mmHg for pH 6.2–7.7 or 17.83 mmHg for pH 8.0–8.3.

Immunoblots

The anti-NBCn1 antibody (provided by Walter Boron, Case Western Reserve University, USA) was generated by immunizing a guinea pig with a GST-fusion protein containing the C-terminal 87 amino-acid residues of rat NBCn1-B (Chen *et al.*, 2007). The characterization of this antigen shows no crossreactivity to other neuronal Na/HCO₃ transporters (Chen *et al.*, 2007). Neurons were scraped in ice-cold homogenization buffer containing 300 mM mannitol, 5 mM HEPES (pH 7.2), 0.1 mg/mL phenylmethanesulphonyl fluoride and 1× protease inhibitor cocktail I (Calbiochem, San Diego, CA, USA).

Cells were homogenized with a 26-gauge needle and centrifuged at 810 g at 4°C for 10 min. Supernatants were collected and assayed for the determination of protein concentration using the Bradford reagents (Sigma–Aldrich, St Louis, MO, USA). Immunoblot analyses were carried out using equal amounts of protein from the samples. Samples were separated on 4–15% sodium dodecylsulfate (SDS) polyacrylamide gels and blotted onto polyvinylidene fluoride (PVDF) membrane. Blots were incubated with the guinea pig anti-NBCn1 antibody in phosphate-buffered saline (PBS) containing 0.05% Tween 20 and 5% nonfat dry milk (dilution 1:500) for 2 h. Blots were washed four times for 10 min and then incubated with horseradish peroxidase (HRP)-conjugated goat antiguinea pig IgG (Millipore, Billerica, MA, USA) for 1 h. Blots were washed and visualized by ECL chemiluminescence (GE Healthcare, Chicago, IL, USA). For β-actin, blots were stripped with 62.6 mM Tris-HCl (pH 6.7), 2% SDS and 0.7% β-mercaptoethanol at 50°C for 10 min, and reprobed with the mouse β-actin antibody (Millipore). Densitometric measurements of NBCn1 and β-actin signals were carried out using IMAGEQUANT image analysis software (Molecular Dynamics, Sunnyvale, CA, USA) according to the manufacturer's protocol. Mean pixel intensities of NBCn1 and β-actin were measured by positioning boxes around protein bands.

Immunofluorescence immunocytochemistry

After enzymatic and mechanical digestion of the hippocampus as described above, neurons (1–2 × 10⁵ cells) were plated on 60-mm dishes containing poly-D-lysine-coated coverslips and incubated for 10–14 days. Cells were fixed with 4% paraformaldehyde, permeabilized with PBS containing 0.1% Triton X-100 (PBT), and blocked in PBT containing 5% normal goat serum (NGS) for 30 min. After washes, coverslips were incubated with the anti-NBCn1 antibody (1:250) in PBS containing 0.025% Triton X-100 and 2.5% NGS at 4°C overnight. Coverslips were rinsed with PBT twice for 20 min and incubated with the goat HRP-conjugated antiguinea pig IgG (1:400; GE Healthcare) for 30 min. Coverslips were then incubated with TSA + tetramethylrhodamine (1:300; Perkin Elmer Life Sciences, Boston, MA, USA). Coverslips were mounted on slides with Vectashield (Vector Laboratories, Burlingame, CA, USA) and examined with a Zeiss Axiovert 100M confocal microscope (Oberkochen, Germany) using a Plan Neofluar 100× lens (numerical aperture 1.3; oil immersion). Immunofluorescence images were acquired using the LSM510 confocal acquisition software. Images were obtained from a single optical section (130 × 130 × 1.49 μm) with a 543-nm fluorescence filter. A single optical section was achieved by stacking images at an interval of 0.5 μm for 12 stacks. Exposure time was 6.29 s for sections. The pinhole setting was 186 μm. Captured images were analyzed using IMAGEQUANT image analysis software. The tetramethylrhodamine immunofluorescence image was converted to a gray scale and inverted. Fluorescence intensity was calibrated by setting the gray value range from 0 (white) to 255 units (black). The fluorescence intensity of tetramethylrhodamine was in the linear range. Mean pixel intensity of immunofluorescence was determined by positioning boxes (700 square pixels) at different locations around the soma and dendrites. Background intensity was determined by positioning boxes in the areas where there was no cell.

Measurements of pH_i

The steady-state pH_i of neurons was determined according to the protocol of Wang *et al.* (2006) with a slight modification. Briefly,

neurons on a coverslip were loaded with $6.5 \mu\text{M}$ of 2',7'-bis-(2-carboxyethyl)-5(6)-carboxyfluorescein acetoxymethyl ester (BCECF-AM) for 10 min and then mounted in a closed-bath confocal imaging chamber RC-30 (Warner Instruments, Hamden, CT, USA) affixed on the stage of a Nikon TE200 inverted microscope. The microscope was equipped with a Lambda 10-2 filter wheel controller and a multi-wavelength filter set. The dye was alternately excited with 500 and 440 nm light, and the emission light at 530 nm was captured. The ratio of the emission from 500 nm (I_{500}) to the emission from 440 nm (I_{440}) was calculated after the subtraction of backgrounds (i.e. neurons without dye) from total I_{500} and I_{440} . Calibration of the fluorescence signal was performed using the K^+/H^+ ionophore nigericin. The steady-state pH_i was calculated by determining linear least-squares analysis over a minimum of 20 s. Data were acquired using METAFLUOR software (Universal Imaging, Downingtown, PA, USA). All experiments were performed at 37°C .

Production of small interference RNA (siRNA)/NBCn1 lentiviral particles

Viral vectors for siRNA/NBCn1-knockdown and siRNA/ns control were made by the custom design and construction service in Cellogenetics (Ijamsville, MD, USA). The nucleotides corresponding to position 589–607, 1641–1659, and 2739–2757 of rat NBCn1 (Genbank accession number: NM_058211) were selected as the best target sequences for siRNA knockdown according to the company's design algorithm. The nucleotides were separately inserted downstream of the human U6 promoter in the lentiviral vector, pLenti-h6BX. The control vector pLenti-hsNS containing 19 non-specific nucleotides GTTCTCCGAACGTGTCACG was purchased from Cellogenetics. The vector contains enhanced green fluorescence protein (eGFP), which is used to detect infection with the green fluorescence filter. Viral particles were produced by transfecting human embryonic kidney HEK293-FT cells with pLenti-h6BX/NBCn1 and pLenti-h6BX/ns vectors. The day before transfection, the HEK293-FT cells (5×10^6 cells) were plated on a 100-mm plate in the culture medium (DMEM with 10% fetal bovine serum, 25 mM HEPES and 120 $\mu\text{g}/\text{mL}$ penicillin and streptomycin, pH 7.4). Next day, the medium was replaced with fresh culture medium without antibiotics. Cells (90–95% confluent) were transfected with 5 μg lentiviral vectors, 2.5 μg pVSV-G (encoding G envelope protein), 3.75 μg pCM78.9 (encoding Gag and Pol proteins) and 36 μL Lipofectamine 2000 (Invitrogen) in 1.5 mL of Opti-MEM medium according to the manufacturer's protocol. Cells were incubated at 37°C overnight, and the medium was replaced with a culture medium containing 2 mM L-glutamine and 0.5 mg/mL Geneticin. The medium was collected 72 h post-transfection and centrifuged at 500 g for 10 min. The supernatant containing viral particles was filtered through a membrane (0.45 μm pore size) and stored at -80°C .

Titration of viral stocks and infection of hippocampal neurons

Titration of viral stocks was done by transducing HEK293 FT cells or HEK cells that stably express NBCn1 (Cooper *et al.*, 2005). The day before titration, cells were plated in a six-well plate ($1.8\text{--}2 \times 10^5$ cells per well) and incubated in the culture medium overnight. On the day of infection, 10-fold serial dilutions (10^{-2} to 10^{-6}) of the viral supernatants were added to 1 mL of fresh culture medium and the medium in the wells was replaced. Cells were incubated at 37°C for 2 days. The infection was monitored by counting green fluorescent cells examined via an inverted microscope Axiovert 135 (Zeiss) with a

GFP filter. The titer in transducing units (TU) was calculated by counting GFP-positive cells in two consecutive diluted plates. The titer was typically 1×10^5 TU/mL or higher. To concentrate the viral stocks, the supernatants were centrifuged at 72,100 g at 16°C for 90 min and pellets were resuspended in PBS and stored at -80°C . The titer of the concentrated viral stocks was typically $\sim 1 \times 10^7$ TU/mL. Hippocampal neurons were then infected with virus at a multiplicity of infection (MOI) of 0.36–1.1. The infection was monitored on a daily basis for green fluorescence. At 72-h post infection, neurons were incubated in the culture medium at pH 6.3 or 7.2 for 6 h and then lysed with the lysis buffer for immunoblot to determine the efficacy of NBCn1 knockdown.

Cell damage and cytotoxicity

Cell damage or cytotoxicity of neurons was determined by the quantitative analysis of lactate dehydrogenase (LDH) released from neurons (Koh & Choi, 1987). Following enzymatic and mechanical digestion of hippocampi, neurons were split in a 96-well plate (1.8×10^4 cells/well) and incubated in Neurobasal medium at 37°C for 10–14 days. Cytotoxicity was induced by treating neurons with 0 or 500 μM glutamate for 10 min in the exposure medium (in mM: NaCl, 119; KCl, 5; CaCl_2 , 2; MgCl_2 , 1.8; glucose, 10; HEPES, 10; glycine, 0.05; and NaHCO_3 , 28; pH 7.2) and then incubating neurons in pH 6.3 or 7.2 media without glutamate for 6 h. MgCl_2 was omitted from Mg^{2+} -free medium. For knockdown experiments, neurons were infected with siRNA/NBCn1 or siRNA/ns lentiviral particles in normal culture medium and then incubated for 72 h before glutamate exposure. Cell damage or cytotoxicity was determined using the LDH-Cytotoxicity Assay kit (BioVision, Mountain View, CA, USA). The maximum amount of LDH release was achieved by breaking neurons with 1% Triton X-100. The media (no cells) with or without Triton X-100 served as backgrounds. The amount of formazan produced in culture supernatants was determined by measuring the absorbance of samples at 490 nm with the microtiter plate reader MRX (Dynex, Chantilly, VA, USA). Cell damage or cytotoxicity was expressed as the percentage of the total LDH release from cell lysis after subtracting backgrounds.

Statistical analysis

The n -value in the manuscript refers to the number of samples that were independently prepared from three to five experiments. Data are reported as mean \pm SE. Level of significance was assessed using: (i) an unpaired, two-tailed Student's t -test to analyze immunofluorescence between neurons at normal vs. acidic pH; (ii) a paired, two-tailed Student's t -test to analyze immunoblot and LDH release of paired groups of neurons (control vs. knockdown) subjected simultaneously to normal and acidic pH; (iii) a one-way ANOVA with the Bonferroni post-test to analyze NBCn1 protein expression and pH_i in a series of pH; and (iv) a two-way ANOVA with the Bonferroni post-test to analyze LDH release from control and knockdown neurons at different time in Mg^{2+} -free medium. $P < 0.05$ was considered significant. Data were analyzed using ORIGIN 8.1 software (OriginLab Corporation; Northampton, MA, USA).

Results

Characterization of the new NBCn1 antibody

We validated the new guinea pig polyclonal antibody against the C-terminal 87 amino acids of rat NBCn1 by immunoblot. These

residues correspond to most of the predicted cytoplasmic C-terminal domain of a rat NBCn1-B splice variant (Choi *et al.*, 2000). The characterization of this antigen shows no cross-reactivity with other neuronal Na/HCO_3^- transporters (Chen *et al.*, 2007). Figure 1A shows an immunoblot of a membrane preparation from the primary cultures of embryonic rat hippocampal neurons. The antibody recognized a band at ~ 150 kDa, similar to NBCn1 protein in cerebral cortex and hippocampus of mouse as detected with a rabbit polyclonal antibody against the same antigenic domain (Chen *et al.*, 2007). The immunoreactive band disappeared when the primary antibody was preabsorbed with the GST/NBCn1 fusion protein containing the C-terminal 87 amino acids of rat NBCn1-B. The specificity of the antibody was also tested by immunoblot of a membrane preparation from *Xenopus* oocytes expressing rat NBCn1 or injected with water (Fig. 1B). The antibody recognized a band at ~ 150 kDa in oocytes expressing the transporter, but not in control oocytes.

NBCn1 in cultured hippocampal neurons was upregulated at low pH

Significant decreases in pH_i and extracellular pH (pH_o) occur during cerebral ischemia (Lipton, 1999). The magnitude of the decrease varies between 0.6 and 1.2 pH units for global and focal ischemia. The decrease occurs within minutes of global ischemia whereas for focal ischemia the decrease occurs within minutes to hours (core region) and hours (penumbra). To examine the effect of acidic pH on NBCn1, we first determined NBCn1 protein expression in primary cultures of rat hippocampal neurons that were incubated in media at pH ranging from 6.2 to 7.4 [established by adjusting (HCO_3^-) at fixed 5% CO_2] for 1 h. Neurons were also incubated in media at pH ranging from 7.4 to 8.3 [established by increasing (HCO_3^-) and lowering P_{CO_2}] to examine the transporter expression at alkaline pH. Crude plasma membranes were isolated from neurons and assessed for immunoblot. Figure 2A shows a marked increase in NBCn1 protein expression at $\text{pH}_o < 6.5$. Changes in NBCn1 expression were negligible at pH_o ranging from 6.8 to 8.3. The signals for β -actin, an internal control, remained unaffected at pH_o ranges we tested. Analyzed by densitometric measurements of NBCn1 relative to β -actin (Fig. 2B), the expression of NBCn1 at $\text{pH}_o < 6.5$ was significantly stronger than those at pH_o




FIG. 1. Characterization of the NBCn1 antibody. (A) Immunoblot of crude plasma membranes isolated from the primary cultures of embryonic rat hippocampal neurons. Immunoblot was performed with the guinea pig NBCn1 antibody against the C-terminal 87 amino acids of rat NBCn1-B. For preabsorption, the antibody was preabsorbed with GST/NBCn1 fusion protein containing the immunogen (50 $\mu\text{g}/\text{mL}$) for 1 h before probing the membrane. (B) Immunoblot of membrane preparation from *Xenopus* oocytes. Oocytes were injected with cRNA encoding rat NBCn1-E or water.




FIG. 2. NBCn1 expression in hippocampal neurons at different pH. (A) Immunoblot of NBCn1. Cultured hippocampal neurons of rats were incubated in media at pH ranging from 6.2 to 8.3 for 1 h. Immunoblot of crude plasma membranes isolated from neurons was performed with an anti-NBCn1 antibody. The blot was striped and reprobed with the anti- β -actin antibody. One of four experiments is shown. (B) Densitometric measurement of NBCn1. The pixel intensity of the NBCn1-immunoreactive band was measured and normalized to that of β -actin ($n = 4$); * $P < 0.05$. (C) Steady-state pH_i of neurons incubated at different pH_o for 1 h. The pH_i was measured using the ratiometric pH-sensitive dye BCECF-AM. Significant decreases in pH_i were observed at $\text{pH}_o < 6.5$. The pH_i values at pH_o 6.5–8.0 were similar ($P > 0.05$, two-way ANOVA). Measurements were made with four to nine preparations at each pH_o ; error bars are SEM.

6.8–8.3 ($F_{7,22} = 6.3$, $P < 0.001$). The expression at pH_o 6.5 was five times higher than that at pH_o 7.4. A similar increase at pH_o 6.2 was also observed when neurons were incubated for 30 min (data not shown).

The steady-state pH_i of neurons at different pH_o was measured using the ratiometric pH-sensitive dye BCECF-AM. Large decreases in pH_i were observed at $\text{pH}_o < 6.5$ ($F_{7,30} = 62.6$, $P < 0.001$; Fig. 2C). At pH_o 6.5, the pH_i was 6.32 ± 0.04 ($n = 8$), significantly lower than 7.55 ± 0.02 ($n = 6$) at pH_o 7.4 ($F_{1,12} = 509.6$, $P < 0.001$). Changes in pH_i at pH_o 6.5–8.0 were relatively small. These data show a close correlation of the pH_i profile with the NBCn1 expression profile: lower pH_i with greater NBCn1 expression.

Figure 3 shows NBCn1 immunocytochemistry of cultured neurons that were incubated at pH_o 6.5 or 7.4 for 1–3 h ($n = 3$ for each). Tetramethylrhodamine-amplified immunofluorescence was localized *in puncta* in somatodendrites, consistent with the previous observation of NBCn1 localization in these neurons (Cooper *et al.*, 2005). Acidic incubation markedly increased NBCn1 immunofluorescence (Fig. 3A and D). The increase occurred in the intensity and number of the punctuate fluorescence in soma and in dendrites at 1 h and remained for 3 h. Controls without the primary antibody had no labeling. Immunofluorescence was quantitated by measuring mean pixel intensity at different locations around single neurons (see Materials and methods). An increase in NBCn1 labeling was detected at low pH_o in membranes ($t_{80} = -20.1$, $P = 0.004$) and cytosol of soma ($t_{60} = -20.7$, $P = 0.005$) as well as in dendrites ($t_{189} = -18.4$, $P < 0.001$; Fig. 3H). Neurons had negligible dendritic blebbing or swelling, indicating that cells were not significantly damaged at pH_o 6.5. The negligible cell damage is compatible with low values of LDH release in these neurons at acidic pH_o as described below.

NBCn1 was upregulated in Mg^{2+} -free medium

In addition to acidification, other ionic changes occur during ischemia and one of these changes is a decrease in the extracellular Mg^{2+} level; for review see McKee *et al.*, 2005). To examine NBCn1 protein expression under these conditions, we incubated cultured hippocampal neurons in Mg^{2+} -free medium and determined NBCn1 expression by immunoblot. Figure 4A shows a representative immunoblot of membrane preparation from neurons that were incubated in Mg^{2+} -free medium at pH 7.2 for 3–6 h. NBCn1 protein expression was upregulated under this condition, while β -actin expression remained unaffected. The densitometric measurements of NBCn1 relative to β -actin showed a 35–40% increase during these incubations ($n = 3$ for each, $t_2 = -5.8$, $P = 0.03$; Fig. 4B). This magnitude was relatively small compared to the fivefold increase at pH_o 6.2–6.5 in Fig. 2, but we obtained consistent results in four independent experiments. The negligible change in NBCn1 expression at 1 h is not shown.

Neurons with NBCn1 knockdown maintained viability

In studying the effect of NBCn1 on cell damage or cytotoxicity, it is not possible to pharmacologically inhibit NBCn1 because no specific blockers are available. For this reason, we developed a lentivirus-mediated siRNA knockdown approach by which NBCn1 transcripts, and subsequent proteins, are degraded in neurons. The nucleotide sequences corresponding to positions 589–607, 1641–1659, and 2739–2757 of rat NBCn1 were selected and separately introduced to a lentiviral vector containing eGFP. The control was a non-specific 19 nucleotide sequence (see Materials and methods). The efficacy of NBCn1 knockdown was tested by infecting HEK cells that stably express the transporter (Cooper *et al.*, 2005) with siRNA/NBCn1 knockdown and siRNA/ns control lentivirus (data not shown). Hippocampal neurons were then infected with knockdown or control lentivirus ($1.8\text{--}5.4 \times 10^5$ viruses per 5×10^5 neurons) and 72 h post-infection the medium was switched to an acidic medium of pH 6.3 for 6 h. Immunoblot showed a marked reduction in NBCn1 expression mediated by siRNA/NBCn1 knockdown (Fig. 5A). Densitometric measurement indicated a reduction of 90% on average ($n = 3$ for each; $t_2 = 11.3$, $P = 0.008$; Fig. 5B).

To examine the effect of NBCn1 knockdown on cell viability, we infected neurons with siRNA/ns or siRNA/NBCn1 lentivirus and 72 h post-infection switched medium to a bath pH of 6.3 or 7.2 for 6 h. pH

6.3 was chosen to ensure NBCn1 upregulation in acidic incubation (i.e. within pH 6.2–6.5). The toxic effect of NBCn1 knockdown was then tested by measuring LDH release from neurons. LDH release from cell lysis represents the total LDH release from neurons treated with 1% Triton X-100, while the release from medium only (no cells) refers to background. Both total and basal releases were unaffected by changes in pH, thus validating the analysis of cell damage at different pH values. Cell damage is presented as the percentage of LDH release normalized to total release after subtracting backgrounds. In neurons with siRNA/ns control (Fig. 5C), the mean cell damage was $5.8 \pm 0.1\%$ at acidic pH incubation and $6.3 \pm 0.7\%$ at normal pH incubation ($n = 6$ for each). These values were not significantly different from each other ($t_5 = -2.5$, $P = 0.06$). Low LDH release at pH_o 6.3 is consistent with the finding by others (Yermolaieva *et al.*, 2004) that cell damage is minimal in cultured hippocampal neurons of mouse at $pH_o > 6.0$. Similarly, in neurons with siRNA/NBCn1 knockdown (Fig. 5D), the mean cell damage was $9.0 \pm 0.5\%$ at acidic pH incubation and $9.2 \pm 0.6\%$ at normal pH incubation ($n = 6$ for each), not significantly different from each other ($t_5 = -2.5$, $P = 0.53$). LDH release from knockdown neurons remained small at both acidic and normal pH. Thus, despite NBCn1 expression being severely reduced by knockdown, neurons maintained their viability at both pH_o 6.3 and 7.2.

NBCn1 knockdown did not affect acidic protection of neurons from glutamate cytotoxicity

The above results indicate that NBCn1 had negligible effect on the viability of neurons maintained at low pH_o although the transporter is upregulated in acidic environments. We then examined whether NBCn1 was linked to acidic protection from cytotoxicity. Cytotoxicity was induced by applying 500 μM glutamate (50 μM glycine and 0 Mg^{2+} , pH_o 7.2) for 10 min (Choi *et al.*, 1987). Neurons were infected with control and knockdown virus and, at 72 h post-infection, neurons were exposed to glutamate and then switched to medium at pH 6.3 or 7.2 for 6 h without the added glutamate. In control neurons (Fig. 6, left bar graphs), glutamate exposure induced cell damage at pH_o 7.2 incubation ($31 \pm 2\%$, $n = 5$). Acidic incubation significantly reduced damage ($7.6 \pm 1.0\%$, $n = 5$; $t_4 = -10.6$, $P < 0.001$). This low value was similar to the basal values (5.8%) for neurons that were incubated at acidic pH without glutamate exposure (Fig. 5A). The decrease following acidic incubation was 75% on average. In NBCn1-knockdown neurons (Fig. 6, right bar graphs), glutamate exposure caused cell damage at pH_o 7.2 ($30 \pm 2\%$, $n = 5$). The damage was significantly reduced at pH_o 6.3 ($6.5 \pm 1.0\%$, $n = 5$; $t_4 = -8.4$, $P = 0.001$), showing a decrease by 78% on average. This decrease was not different from the decrease for control neurons ($F_{1,17} = 1.1$, $P = 0.31$). Thus, neurons were protected from glutamate cytotoxicity by acidic pH_o regardless of NBCn1 knockdown.

NBCn1 knockdown reduced cytotoxicity in a prolonged Mg^{2+} -free incubation

NBCn1 was upregulated in Mg^{2+} -free medium (Fig. 4). To examine whether this upregulation was linked to cytotoxicity, we exposed neurons to glutamate as described above and incubated them in Mg^{2+} -free medium for 1–6 h (pH 7.2). Figure 7A shows that glutamate cytotoxicity progressively increased with time in both control neurons and NBCn1-knockdown neurons. This development of cytotoxicity is consistent with the previous report (Sun *et al.*,




FIG. 3. Immunocytochemistry of NBCn1 in hippocampal neurons. (A–E) NBCn1 immunofluorescence in neurons incubated at pH 6.5 or 7.4 for 1–3 h. The labeling was amplified with tetramethylrhodamine (see Materials and methods). Control was neurons without the primary antibody. One of three immunofluorescence experiments is shown. (F and G) NBCn1 immunofluorescence in dendrites of neurons incubated at pH 6.5 or 7.4 for 1 h. The whole cell images are shown in insets. (H) Quantitation of NBCn1 immunofluorescence. The immunofluorescence images were converted to a gray scale and inverted (0 units as white and 255 units as black). Mean pixel intensity of NBCn1 immunofluorescence was determined by positioning boxes (700 square pixels) at different locations around plasma membrane, cytosol and dendrites. Numbers in parentheses are numbers of counts, and error bars are SEM. Scale bars, 50 μ m. * P < 0.05.




FIG. 4. NBCn1 expression in Mg^{2+} -free medium. (A) Immunoblot of NBCn1. Cultured hippocampal neurons were incubated in Mg^{2+} -free medium (pH 7.2) for 3–6 h. Immunoblots of crude plasma membrane preparation from neurons were probed with the anti-NBCn1 antibody. The blot was striped and reprobed with the anti- β -actin antibody. (B) Densitometric measurements of NBCn1. The pixel intensity of NBCn1 was measured and normalized to that of β -actin ($n = 4$ for each). Error bars are SEM. * $P < 0.05$.

2001) that a prolonged Mg^{2+} depletion induces spontaneous epileptiform-like activity and causes cell death in hippocampal neurons. However, glutamate cytotoxicity in a prolonged Mg^{2+} -free incubation was attenuated in NBCn1-knockdown neurons ($F_{1,54} = 11.9$, $P = 0.001$). The difference between the two groups of neurons was progressively more prominent with time. Thus, at 6 h incubation, NBCn1-knockdown neurons had ~30% less cytotoxicity than controls. A similar Mg^{2+} -free incubation without glutamate exposure had negligible cytotoxicity in both groups of neurons (Fig. 7B), thus indicating that Mg^{2+} -free incubation itself is not sufficient to cause cytotoxicity. To test whether the reduction in cytotoxicity by NBCn1 knockdown was dependent upon pH, we incubated control neurons and NBCn1-knockdown neurons in acidic medium without Mg^{2+} after glutamate exposure. However, this approach did not provide valuable information as both groups of neurons had a reduction in cytotoxicity due to the protective effect of acidity (data not shown). As an alternative approach, we incubated neurons at pH 6.3 for 1 h before glutamate exposure (Fig. 7C). Following incubation at acidic pH, NBCn1-knockdown neurons had lower glutamate cytotoxicity than control neurons ($t_9 = 4.0$, $P = 0.003$). The decrease was ~16% on average, not significantly different from 23% glutamate cytotoxicity on average at pH_o 7.2 preincubation ($F_{1,29} = 2.7$, $P = 0.11$). Therefore, NBCn1 knockdown reduced glutamate cytotoxicity regardless of acidic preconditioning.

In conclusion, our data show that NBCn1 in cultured hippocampal neurons was upregulated at low pH and in Mg^{2+} -free medium. The




FIG. 5. Effect of NBCn1 knockdown on viability of neurons. (A) Immunoblots of NBCn1 in neurons treated with control and knockdown lentivirus. (B) Densitometric measurements of NBCn1. The intensity of NBCn1 was normalized to that of β -actin ($n = 3$). (C) LDH release from neurons infected with siRNA/ns control lentivirus. Neurons were incubated in media at pH 6.3 or 7.2 for 6 h and LDH release was determined by measuring the medium absorbance at 490 nm. The absorbance value from cell lysis represents the total LDH release from neurons lysed with 1% Triton X-100, and the value from medium is the background without cells. (D) LDH release from neurons infected with siRNA/NBCn1 knockdown lentivirus ($n = 6$). Error bars are SEM. * $P < 0.05$.

upregulation had minimal effect on acidic protection from glutamate cytotoxicity. In the absence of Mg^{2+} , however, the upregulation could make neurons more vulnerable to glutamate cytotoxicity. The significance of our findings is discussed below.

Discussion

In this study, we provide the first evidence for the association of NBCn1 with cytotoxicity in cultured hippocampal neurons. Using




FIG. 6. Glutamate cytotoxicity in neurons at acidic and normal pH. Neurons with siRNA control and NBCn1 knockdown lentivirus (72 h post-infection) were exposed to 500 μ M glutamate (50 μ M glycine and 0 Mg^{2+} , pH 7.2) for 10 min and then switched to medium at pH 6.3 or 7.2 for 6 h without the added glutamate. Cytotoxicity was determined as the percentage of LDH released from neurons relative to the total release after background subtraction ($n = 5$ for each groups of neurons). Error bars are SEM. * $P < 0.05$.

molecular and cellular approaches, we determined the effects of acidity and Mg^{2+} on NBCn1 expression, knocked down endogenous NBCn1 expression in neurons, and examined the effect of NBCn1 knockdown on glutamate cytotoxicity under conditions where the transporter was upregulated. The major finding is that NBCn1 in hippocampal neurons had a dynamic upregulation in low pH as well as Mg^{2+} -free incubation, but its contribution to glutamate cytotoxicity varied depending upon the availability of acidity and Mg^{2+} . Our data could be a valuable starting point for future studies aimed at determining the pathological role of NBCn1 in cerebral ischemia.

Upregulation of NBCn1

A dramatic increase in NBCn1 protein expression occurred when neurons were incubated at pH 6.2–6.5 (Fig. 2). The expression remained unaffected at pH_o 6.5–8.3, which is in the range of ~ 1 pH unit from the physiological pH 7.2–7.4. NBCn1 is known to have a dynamic response to acidic environments in some tissues. In the kidney, the expression and activity of NBCn1 is upregulated by chronic metabolic acidosis (Kwon *et al.*, 2002; Jakobsen *et al.*, 2004). Chronic metabolic acidosis increases NH_4^+ absorption from the lumen in the medullary thick ascending limb of the kidney. The increase in NH_4^+ absorption overloads intracellular H^+ in the tubules as NH_4^+ dissociates into NH_3 and H^+ (Knepper *et al.*, 1989). The thick ascending limb cells compensate intracellular H^+ overloads by upregulating NBCn1 in chronic metabolic acidosis. Whether a similar NBCn1-dependent compensatory mechanism exists in neurons is unclear, but our study shows a close correlation between low pH_i and NBCn1 upregulation in primary cultures of hippocampal neurons. The pH_i of cultured neurons markedly decreased at $\text{pH}_o < 6.5$ (Fig. 2C) and this decrease was similar to NBCn1 upregulation at these pH_o ranges. In adjusting pH_o , we varied (HCO_3^-) and P_{CO_2} in the medium. The (HCO_3^-) in the medium of pH 6.5 is eight times less than that at pH_o 7.4 (3.25 vs. 26 mM). Despite this low (HCO_3^-) in the medium, NBCn1 was markedly upregulated at pH_o 6.5. Conversely, the (HCO_3^-) in the medium of pH 8.3 was four times higher (104 vs. 26 mM). Despite this, NBCn1 expression remained unaffected at pH_o 8.3. Therefore,

it is unlikely that HCO_3^- in the medium determined NBCn1 expression.

Negligible role of NBCn1 in acidic protection

With the observation of NBCn1 upregulation in acidic environments, we hypothesized that the transporter would play a role in protecting neurons from cytotoxicity at low pH. We instead found both control neurons and NBCn1-knockdown neurons having a similar reduction in glutamate toxicity at low pH_o (Fig. 6), thus indicating a negligible effect of NBCn1 knockdown on acidic protection. In our experiments, the glutamate exposure condition was identical in both control neurons and knockdown neurons before they were placed in either acidic or normal medium. This suggests that the protection of neurons at acidic pH_o is not caused by the process of glutamate exposure but instead is directly related to the acidity of media or in the cell. It is not clear how the acidity of media or in the cell reduces cytotoxicity mediated by glutamate exposure, but the mechanism should involve inhibition of a pH-dependent process downstream of glutamate receptor activation.

Effect of NBCn1 on glutamate cytotoxicity caused by Mg^{2+} -free incubation

Despite the negligible role of NBCn1 in acidic protection, the transporter can affect neuronal damage. NBCn1-knockdown neurons had an attenuation of glutamate cytotoxicity in a prolonged Mg^{2+} -free incubation (Fig. 7A). This attenuation did not occur when cytotoxicity was examined in the presence of Mg^{2+} (Fig. 6), thus indicating that the effect of NBCn1 on neuronal damage was dependent upon the availability of Mg^{2+} in the medium. In our experiments, ' Mg^{2+} -free' refers to the nominal removal of Mg^{2+} from the medium but does not imply a complete lack of Mg^{2+} , as discussed by others (Sombati & DeLorenzo, 1995). Mg^{2+} may be released from the cell via Mg^{2+} transporters (Schmitz *et al.*, 2007). The exact concentration of Mg^{2+} in our Mg^{2+} -free medium is not determined, but it would probably be lower than that *in vivo* during pathological states. The Mg^{2+} level is reduced to 0.3–0.48 mM within 8 h after the onset of ischemia in human (Altura *et al.*, 1997). It remains to be determined whether the effect of NBCn1 on damage of hippocampal neurons can occur at such Mg^{2+} levels.

By what mechanism does NBCn1 affect cytotoxicity caused by Mg^{2+} depletion? The development of cytotoxicity requires glutamate (Fig. 7B) and this led us to speculate that Mg^{2+} -sensitive N-methyl-D-aspartate (NMDA) receptors may be involved in cytotoxicity under Mg^{2+} -free conditions. The receptors mediate a large intracellular Ca^{2+} load that initiates apoptotic cellular events leading to cell death (Dingledine *et al.*, 1999). The receptors are blocked by Mg^{2+} in a voltage-sensitive and noncompetitive manner. Taken together, it is possible that Mg^{2+} depletion from the glutamate-containing medium causes NMDA receptor activation, which results in cytotoxicity.

Based on the data in the present study, we propose a model for the pathological effect of NBCn1 on cytotoxicity in hippocampal neurons. The pH_i of neurons falls under acidic conditions and the acid-extruder NBCn1 is upregulated to compensate for intracellular acidification. The upregulation of NBCn1 may cause neuronal damage, but this does not occur at low pH_i due to the protective effect of acidity. The transporter can also be upregulated at normal pH_o when Mg^{2+} is less available. Under this low- Mg^{2+} condition, the upregulation of the transporter facilitates deleterious effects on neurons and renders




FIG. 7. Effects of NBCn1 knockdown on cytotoxicity in Mg^{2+} -free medium. (A) Glutamate cytotoxicity in Mg^{2+} -free medium. Neurons with siRNA control and knockdown were incubated for 1–6 h in Mg^{2+} -free medium (pH 7.2) after glutamate exposure and assessed for the cytotoxicity assay ($n = 5$ for each). (B) Cytotoxicity in Mg^{2+} -free medium without glutamate exposure. The experimental procedure was similar to the above except that neurons were exposed to 0 mM glutamate ($n = 5$ for each). (C) Preincubation of neurons at pH 6.3 before glutamate exposure. Neurons were incubated at pH 6.3 or 7.2 before glutamate exposure and then incubated in Mg^{2+} -free medium (pH 7.2) for 2.5 h ($n = 10$ for pH 6.3 and $n = 5$ for pH 7.2). Error bars are SEM. * $P < 0.05$.

neurons more vulnerable to cytotoxicity. Thus, the cytotoxicity potentially mediated by the transporter is normally protected but can be uncovered by low Mg^{2+} in the extracellular fluid of the brain. NBCn1 could be one of the proteins to facilitate neuronal injury in hypomagnesemia.

Comparison with Na/HCO_3 transporter-mediated damage in astrocytes

The electrogenic Na/HCO_3 cotransporter (NBCe1; SLC4A4) is predominantly expressed in astrocytes (Rickmann *et al.*, 2007; Majumdar *et al.*, 2008). Inhibition of this transporter reduces cell vulnerability to acid injury (Giffard *et al.*, 2000). The data in our study are thus consistent with the report by Giffard *et al.* (2000) in that inhibition or knockdown of Na/HCO_3 transporters reduces cell damage. Despite this similarity, several differences are recognizable. First, hippocampal neurons are viable at pH 6.3 incubation whereas astrocytes die at pH 6.8 incubation. This difference is consistent with the previous reports that neurons are more resistant than astrocytes to acidic injury. Second, NBCn1 knockdown causes negligible effects on neuronal viability at low pH_o whereas NBCe1 inhibition increases glial viability at low pH_o . Third, NBCn1 contributes to glutamate cytotoxicity in Mg^{2+} -free conditions whereas NBCe1 appears to have a limited effect on acidic injury. Apparently, both NBCn1 and NBCe1 affect cytotox-

icity or cell death, but the mechanisms underlying their effects are distinct.

Maintenance of cell viability at low pH

In addition to the finding of a pathological role of NBCn1, we also observed that cultured hippocampal neurons maintained their viability at pH_o 6.3 (Fig. 5). The threshold pH_o for acidic cell death depends on both time and degree of intracellular acidification. In cortical cultures of neurons, the threshold to induce 50% cell death is pH_o 6.4 for 6 h acid exposure in the absence of CO_2/HCO_3^- buffer (Nedergaard *et al.*, 1991). In cultured hippocampal neurons of mouse, LDH release at $pH_o > 6.0$ is minimal (Yermolaieva *et al.*, 2004). A prolong exposure to an acidic environment can induce the acid-activated ion channels (ASIC) in central neurons (Johnson *et al.*, 2001; Xiong *et al.*, 2004), and mediates Ca^{2+} influx via ASIC, causing acidic injury in neurons. However, these channels are mostly activated at $pH_o < 6.0$ (Johnson *et al.*, 2001; Xiong *et al.*, 2004). Consistent with this, our data show low LDH release from hippocampal neurons at pH_o 6.3.

Summary

We found that NBCn1 knockdown reduces cytotoxicity of hippocampal neurons subjected to Mg^{2+} depletion from the medium. This

finding is consistent with the previous reports that inhibition of Na/H exchange or Na/HCO₃ cotransport protects cells against damage or injury in cardiac cells. Nevertheless, in our system, the mechanism for NBCn1 contribution to cytotoxicity appears to be different from those in cardiac cells, and further investigation is needed. In particular, it will be interesting to investigate what mechanisms ameliorate the cytotoxicity that would be predicted with the upregulation of NBCn1.

Acknowledgements

The authors acknowledge Walter Boron (Case Western Reserve) for providing the guinea pig polyclonal NBCn1 antibody. The authors acknowledge Ronald Abercrombie (Emory) for helpful discussion. This work was supported by the American Heart Association Southeast Affiliate and the URC grant at Emory (I.C.), and NIH DK61418 (C.C.Y.).

Abbreviations

BCECF-AM, 2',7'-bis-(2-carboxyethyl)-5(6)-carboxyfluorescein acetoxymethyl ester; DMEM, Dulbecco's modification of Eagle's medium; eGFP, enhanced green fluorescence protein; HEPES, 4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid; HRP, horseradish peroxidase; LDH, lactate dehydrogenase; MOI, multiplicity of infection; NBCe1, electrogenic Na/bicarbonate cotransporter 1; NBCn1, electroneutral Na/bicarbonate cotransporter 1; NGS, normal goat serum; PBS, phosphate-buffered saline; pHi, intracellular pH; pHo, extracellular pH; PVDF, polyvinylidene fluoride; siRNA, small interference RNA; SLC4A, solute carrier family transporter 4A; TU, transducing unit.

References

- Altura, B.T., Memon, Z.I., Zhang, A., Cheng, T.P., Silverman, R., Cracco, R.Q. & Altura, B.M. (1997) Low levels of serum ionized magnesium are found in patients early after stroke which result in rapid elevation in cytosolic free calcium and spasm in cerebral vascular muscle cells. *Neurosci. Lett.*, **230**, 37–40.
- Baxter, K.A. & Church, J. (1996) Characterization of acid extrusion mechanisms in cultured fetal rat hippocampal neurones. *J. Physiol.*, **493**, 457–470.
- Bobulescu, I.A., Di Sole, F. & Moe, O.W. (2005) Na⁺/H⁺ exchangers: physiology and link to hypertension and organ ischemia. *Curr. Opin. Nephrol. Hypertens.*, **14**, 485–494.
- Boedtker, E., Praetorius, J., Fuchtbauer, E.M. & Aalkjaer, C. (2008) Antibody-independent localization of the electroneutral Na⁺-HCO₃⁻ cotransporter NBCn1 (slc4a7) in mice. *Am. J. Physiol. Cell. Physiol.*, **294**, C591–C603.
- Chen, L.M., Choi, I., Haddad, G.G. & Boron, W.F. (2007) Chronic continuous hypoxia decreases the expression of SLC4A7 (NBCn1) and SLC4A10 (NCBE) in mouse brain. *Am. J. Physiol. Regul. Integr. Comp. Physiol.*, **293**, R2412–R2420.
- Chen, L.M., Kelly, M.L., Rojas, J.d., Parker, M.D., Gill, H.S., Davis, B.A. & Boron, W.F. (2008) Use of a new polyclonal antibody to study the distribution and glycosylation of the sodium-coupled bicarbonate transporter NCBE in rodent brain. *Neuroscience*, **151**, 374–385.
- Chesler, M. (2003) Regulation and modulation of pH in the brain. *Physiol. Rev.*, **83**, 1183–1221.
- Choi, D.W., Maulucci-Gedde, M. & Kriegstein, A.R. (1987) Glutamate neurotoxicity in cortical cell culture. *J. Neurosci.*, **7**, 357–368.
- Choi, I., Aalkjaer, C., Boulpaep, E.L. & Boron, W.F. (2000) An electroneutral sodium/bicarbonate cotransporter NBCn1 and associated sodium channel. *Nature*, **405**, 571–575.
- Cooper, D.S., Saxena, N.C., Yang, H.S., Lee, H.J., Moring, A.G., Lee, A. & Choi, I. (2005) Molecular and functional characterization of the electroneutral Na/HCO₃ cotransporter NBCn1 in rat hippocampal neurons. *J. Biol. Chem.*, **280**, 17823–17830.
- Dingledine, R., Borges, K., Bowie, D. & Traynelis, S.F. (1999) The glutamate receptor ion channels. *Pharmacol. Rev.*, **51**, 7–61.
- Giffard, R.G., Papadopoulos, M.C., van Hoof, J.A., Xu, L., Giuffrida, R. & Monyer, H. (2000) The electrogenic sodium bicarbonate cotransporter: developmental expression in rat brain and possible role in acid vulnerability. *J. Neurosci.*, **20**, 1001–1008.
- Giffard, R.G., Lee, Y.S., Ouyang, Y.B., Murphy, S.L. & Monyer, H. (2003) Two variants of the rat brain sodium-driven chloride bicarbonate exchanger (NCBE): developmental expression and addition of a PDZ motif. *Eur. J. Neurosci.*, **18**, 2935–2945.
- Grichtchenko, I. I., Risso-Bradley, S., Rojas, J. d., Richerson, G. B. & Boron, W. F. (2001) Localization of the Na⁺-driven Cl-HCO₃ exchanger protein (NDCBE1) in rat brain. *Soc. Neurosci. Abstr.*, A527.12.
- Jakobsen, J.K., Odgaard, E., Wang, W., Elkjaer, M.L., Nielsen, S., Aalkjaer, C. & Leipziger, J. (2004) Functional up-regulation of basolateral Na⁺-dependent HCO₃⁻ transporter NBCn1 in medullary thick ascending limb of K⁺-depleted rats. *Pflügers Arch.*, **448**, 571–578.
- Johnson, M.B., Jin, K., Minami, M., Chen, D. & Simon, R.P. (2001) Global ischemia induces expression of acid-sensing ion channel 2a in rat brain. *J. Cereb. Blood Flow Metab.*, **21**, 734–740.
- Karmazyn, M. (1999) The role of the myocardial sodium-hydrogen exchanger in mediating ischemic and reperfusion injury. From amiloride to cariporide. *Ann. N.Y. Acad. Sci.*, **874**, 326–334.
- Khandoudi, N., Albadine, J., Robert, P., Krief, S., Berrebi-Bertrand, I., Martin, X., Bevensee, M.O., Boron, W.F. & Bril, A. (2001) Inhibition of the cardiac electrogenic sodium bicarbonate cotransporter reduces ischemic injury. *Cardiovasc. Res.*, **52**, 387–396.
- Knepper, M.A., Packer, R. & Good, D.W. (1989) Ammonium transport in the kidney. *Physiol. Rev.*, **69**, 179–249.
- Koh, J.Y. & Choi, D.W. (1987) Quantitative determination of glutamate mediated cortical neuronal injury in cell culture by lactate dehydrogenase efflux assay. *J. Neurosci. Methods*, **20**, 83–90.
- Kwon, T.H., Fulton, C., Wang, W., Kurtz, I., Frokiaer, J., Aalkjaer, C. & Nielsen, S. (2002) Chronic metabolic acidosis upregulates rat kidney Na-HCO₃ cotransporters NBCn1 and NBC3 but not NBC1. *Am. J. Physiol. Renal. Physiol.*, **282**, F341–F351.
- Lipton, P. (1999) Ischemic cell death in brain neurons. *Physiol. Rev.*, **79**, 1431–1568.
- Majumdar, D., Maunsbach, A.B., Shacka, J.J., Williams, J.B., Berger, U.V., Schultz, K.P., Harkins, L.E., Boron, W.F., Roth, K.A. & Bevensee, M.O. (2008) Localization of electrogenic Na/bicarbonate cotransporter NBCe1 variants in rat brain. *Neuroscience*, **155**, 818–832.
- McKee, J.A., Brewer, R.P., Macy, G.E., Borel, C.O., Reynolds, J.D. & Warner, D.S. (2005) Magnesium neuroprotection is limited in humans with acute brain injury. *Neurocrit. Care.*, **2**, 342–351.
- Nedergaard, M., Goldman, S.A., Desai, S. & Pulsinelli, W.A. (1991) Acid-induced death in neurons and glia. *J. Neurosci.*, **11**, 2489–2497.
- Rickmann, M., Orłowski, B., Heupel, K. & Roussa, E. (2007) Distinct expression and subcellular localization patterns of Na⁺/HCO₃⁻ cotransporter (SLC 4A4) variants NBCe1-A and NBCe1-B in mouse brain. *Neuroscience*, **146**, 1220–1231.
- Romero, M.F., Fulton, C.M. & Boron, W.F. (2004) The SLC4 family of HCO₃⁻ transporters. *Pflügers Arch.*, **447**, 495–509.
- Schmitz, C., Deason, F. & Perraud, A.L. (2007) Molecular components of vertebrate Mg²⁺-homeostasis regulation. *Magn. Res.*, **20**, 6–18.
- Schwiening, C.J. & Boron, W.F. (1994) Regulation of intracellular pH in pyramidal neurons from the rat hippocampus by Na⁺-dependent Cl⁻-HCO₃⁻ exchange. *J. Physiol.*, **475**, 59–67.
- Sombati, S. & DeLorenzo, R.J. (1995) Recurrent spontaneous seizure activity in hippocampal neuronal networks in culture. *J. Neurophysiol.*, **73**, 1706–1711.
- Sun, D.A., Sombati, S. & DeLorenzo, R.J. (2001) Glutamate injury-induced epileptogenesis in hippocampal neurons: an in vitro model of stroke-induced "epilepsy". *Stroke*, **32**, 2344–2350.
- Wang, Y., Meyer, J.W., Ashraf, M. & Shull, G.E. (2003) Mice with a null mutation in the NHE1 Na⁺-H⁺ exchanger are resistant to cardiac ischemia-reperfusion injury. *Circ. Res.*, **93**, 776–782.
- Wang, D., Lee, H.J., Cooper, D.S., Cebotaro, L., Walden, P.D., Choi, I. & Yun, C.C. (2006) Coexpression of MAST205 inhibits the activity of Na⁺/H⁺ exchanger NHE3. *Am. J. Physiol. Renal. Physiol.*, **290**, F428–F437.
- Xiong, Z.G., Zhu, X.M., Chu, X.P., Minami, M., Hey, J., Wei, W.L., MacDonald, J.F., Wemmie, J.A., Price, M.P., Welsh, M.J. & Simon, R.P. (2004) Neuroprotection in ischemia: blocking calcium-permeable acid-sensing ion channels. *Cell*, **118**, 687–698.
- Yermolaieva, O., Leonard, A.S., Schnizler, M.K., Abboud, F.M. & Welsh, M.J. (2004) Extracellular acidosis increases neuronal cell calcium by activating acid-sensing ion channel 1a. *Proc. Natl Acad. Sci. USA*, **101**, 6752–6757.