

Based on the alluring structures of cyclopropyl nucleosides and the antiviral activities shown by acyclic phosphonic acid nucleosides, we were inspired to synthesize novel classes of methyl branched cyclopropyl phosphonic acids as potential antiviral agents.

MATERIALS AND METHODS

The melting points were determined on a Mel-tem II laboratory device and were uncorrected. The NMR spectra were recorded on a JEOL JNM-LA 300 spectrometer. The chemical shifts are reported as parts per million (δ) and the signals are quoted as s (singlet), d (doublet), t (triplet), q (quartet), m (multiplet) and dd (doublet of doublets). The UV spectra were obtained using a Beckman DU-7 spectrophotometer. Thin Layer Chromatography (TLC) was performed on Uniplates (silica gel) which were purchased from Analtech Co. All reactions were carried out under N_2 unless otherwise specified. Dry dichloromethane, benzene, and pyridine were obtained by distillation from CaH_2 . Dry THF was obtained by distillation from Na and benzophenone, immediately prior to use.

(\pm)-*trans*-[2-(*tert*-Butyl-dimethyl-silyloxymethyl)-2-methyl-cyclopropyl]-methanol (**8**)

To a mixture comprising of allylic alcohol (**7**) (1.87 g, 8.64 mmol) in 30 mL of CH_2Cl_2 , $ZnEt_2$ (17.3 mL, 1 M in hexane) and CH_2I_2 (9.26 g, 34.58 mmol) were added at 0 °C. The mixture was stirred for 4 h at the same temperature and was quenched by the addition of a saturated solution of NH_4Cl . After the mixture was concentrated to 1/3 of its original volume, the aqueous layer was extracted with EtOAc. The organic layer was dried over anhydrous $MgSO_4$, filtered and evaporated *in vacuo*. The residue was purified by silica gel column chromatography (EtOAc/hexane, 1:12) to form **8** (1.61 g, 81%) as a colorless oil: 1H -NMR ($CDCl_3$, 300 MHz) δ 3.69 (dd, $J = 9.6, 5.1$ Hz, 1H), 3.47 (dd, $J = 10.8, 8.7$ Hz, 1H), 3.39 (d, $J = 9.9$ Hz, 1H), 3.27 (d, $J = 9.9$ Hz, 1H), 1.10 (s, 3H), 0.99 (m, 1H), 0.85 (s, 9H), 0.61 (m, 1H), 0.15 (m, 1H), 0.11 (s, 6H); ^{13}C -NMR ($CDCl_3$) δ 70.54, 63.43, 25.94, 23.30, 22.12, 18.36, 15.33, 14.82, -5.30.

(\pm)-*trans*-[2-(*tert*-Butyldimethylsilyloxymethyl)-2-methyl-cyclopropylmethoxymethyl]-phosphonic acid diisopropyl ester (**9**)

To a solution of **8** (1.64 g, 7.11 mmol) in 6 mL of DMF, $LiOt$ -Bu (72 mg, 0.53 mmol) was added at 25 °C. $LiOt$ -Bu (11.4 mL of 1.0 M solution in THF, 11.4 mmol) and a solution of diisopropyl bromomethylphosphonate (2.5 g, 9.63 mmol) in 6 mL of DMF were slowly and simultaneously added to the reaction mixture for 4 h at 60 °C under anhydrous

condition. The mixture was quenched by adding water (35 mL), and the organic solvents (THF) were removed *in vacuo*. The aqueous layer was extracted with EtOAc (3 \times 60 mL). The combined extracts were washed with brine, dried over $MgSO_4$, and concentrated *in vacuo*. The residue was purified by silica gel column chromatography (EtOAc/hexane, 1:2) to form **9** (1.89 g, 65%) as a colorless syrup: 1H -NMR ($CDCl_3$, 300 MHz) δ 4.75 (m, 2H), 3.71 (d, $J = 8.0$ Hz, 2H), 3.68 (m, 2H), 3.40 (d, $J = 9.8$ Hz, 1H), 3.28 (d, $J = 9.8$ Hz, 1H), 1.31 (m, 12H), 1.09 (s, 3H), 1.00 (m, 1H), 0.82 (s, 9H), 0.56 (m, 1H), 0.21 (m, 1H), 0.12 (s, 6H); ^{13}C -NMR ($CDCl_3$) δ 71.71, 68.31, 66.54, 64.21, 25.67, 23.99, 23.45, 22.11, 18.45, 15.78, 14.65, -5.45; Anal calc for $C_{19}H_{41}O_5PSi$: C, 55.85; H, 10.11. Found: C, 56.02; H, 10.22.

(\pm)-*trans*-[2-(Hydroxymethyl)-2-methyl-cyclopropyl-methoxymethyl]-phosphonic acid diisopropyl ester (**10**)

To a solution of **9** (3.1 g, 7.60 mmol) in tetrahydrofuran (20 mL), tetrabutylammonium fluoride (9.88 mL, 1.0 M solution in THF) was added at 0 °C and stirred for 5 h at room temperature. The reaction mixture was concentrated *in vacuo* and the residue was purified by silica gel column chromatography (EtOAc/hexane, 4:1) to form **10** (1.85 g, 83%) as a colorless syrup: 1H -NMR ($CDCl_3$, 300 MHz) δ 4.70 (m, 2H), 3.73 (d, $J = 8.2$ Hz, 2H), 3.71 (m, 2H), 3.41 (m, 2H), 1.33 (m, 12H), 1.12 (s, 3H), 1.07 (m, 1H), 0.60 (m, 1H), 0.23 (m, 1H); ^{13}C -NMR ($CDCl_3$) δ 70.89, 69.45, 67.33, 65.12, 23.90, 23.23, 21.76, 14.88, 14.01.

(\pm)-*trans*-[Methanesulfonic acid-2-(diisopropoxy-phosphorylmethoxymethyl)-2-methyl-cyclopropyl-methyl] ester (**11**)

To a solution of phosphonate **10** (1.37 g, 4.68 mmol) in anhydrous CH_2Cl_2 (2 mL), anhydrous triethylamine (1.5 mL) and $MsCl$ (642 mg, 5.61 mmol) were added at 0 °C. The mixture was stirred at the same temperature for 5 h, and was quenched by the addition of cold saturated $NaHCO_3$ solution (2.0 mL). The mixture was extracted twice with CH_2Cl_2 (120 mL)/water (120 mL). The combined organic layer was dried over anhydrous magnesium sulfate and filtered. The filtrate was concentrated *in vacuo*, and the residue was purified by silica gel column chromatography (EtOAc/hexane, 4:1) to form **11** (1.16 g, 67%) as a colorless syrup: 1H -NMR ($CDCl_3$, 300 MHz) δ 4.71 (m, 2H), 3.75 (d, $J = 7.8$ Hz, 2H), 3.70-3.66 (m, 2H), 3.44 (dd, $J = 13.5, 8.8$ Hz, 2H), 3.04 (s, 3H), 1.33 (m, 12H), 1.04 (s, 3H), 1.01 (m, 1H), 0.61 (m, 1H), 0.26 (m, 1H); ^{13}C -NMR ($CDCl_3$) δ 70.67, 69.12, 68.56, 65.21, 36.76, 24.21, 23.67, 22.17, 15.67, 14.32.

(±)-trans-9-[2-(Diisopropoxy-phosphorylmethoxymethyl)-2-methyl-cyclopropylmethylester]-adenine (12)

Mixture of mesylate **11** (311 mg, 0.837 mmol), K₂CO₃ (231 mg, 1.75 mmol), 18-crown-6 (221 mg, 0.837 mmol), and adenine (135.8 mg, 1.00 mmol) in dry DMF (9.0 mL) was stirred overnight at 90°C. The mixture was cooled to room temperature and concentrated *in high vacuo*. The residue was diluted with brine (30 mL) and extracted with CH₂Cl₂ (50 mL×3). The combined organic layer was dried over anhydrous MgSO₄, filtered, and concentrated under reduced pressure. The residue was purified by silica gel column chromatography (MeOH/CH₂Cl₂, 1:10) to form compound **12** (130.8 mg, 38%) as a yellowish solid: ¹H-NMR (CDCl₃, 300 MHz) δ 8.45 (s, 1H), 8.02 (s, 1H), 4.72 (m, 2H), 4.61 (dd, *J* = 14.7, 6.6 Hz, 1H), 4.22 (dd, *J* = 14.4, 8.7 Hz, 1H), 3.71 (d, *J* = 8.4 Hz, 2H), 3.61 (d, *J* = 10.2 Hz, 1H), 3.22 (d, *J* = 10.2 Hz, 1H), 1.30 (m, 12H), 1.11 (s, 3H), 1.02 (m, 1H), 0.59 (m, 1H), 0.39 (t, *J* = 5.1 Hz, 1H); ¹³C-NMR (CDCl₃) δ 154.28, 152.65, 141.54, 118.76, 71.31, 70.22, 65.77, 50.61, 23.45, 23.15, 18.33, 15.66, 15.59.

(±)-trans-1-[2-(Diisopropoxy-phosphorylmethoxymethyl)-2-methyl-cyclopropylmethyl ester]-cytosine (13)

Compound **13** was prepared from **11** by following the method as described for **12**: the yield of **13** was 31%: ¹H-NMR (CDCl₃, 300 MHz) δ 7.66 (d, *J* = 7.5 Hz, 1H), 6.95 (br d, 2H), 5.63 (d, *J* = 7.5 Hz, 1H), 4.75 (m, 2H), 3.71-3.53 (m, 4H), 3.48 (d, *J* = 10.2 Hz, 1H), 3.32 (d, *J* = 10.2 Hz, 1H), 1.35 (m, 12H), 1.09 (s, 3H), 0.52 (m, 1H), 0.26 (m, 1H); ¹³C-NMR (CDCl₃) δ 165.74, 155.92, 145.01, 92.91, 70.35, 69.48, 66.35, 48.02, 23.23, 22.01, 17.93, 15.26, 14.35.

(±)-trans-1-[2-(Diisopropoxy-phosphorylmethoxymethyl)-cyclopropylmethyl ester]-thymine (14)

Compound **14** was prepared from **11** by following the method as described for **12**: the yield of **14** was 33%: ¹H-NMR (CDCl₃, 300 MHz) δ 7.87 (s, 1H), 4.79 (m, 2H), 4.01 (dd, *J* = 13.5, 8.6 Hz, 1H), 3.74 (d, *J* = 7.8 Hz, 2H), 3.68 (d, *J* = 10.0 Hz, 1H), 3.47 (dd, *J* = 14.0, 9.9 Hz, 1H), 3.11 (d, *J* = 11.1 Hz, 1H), 1.97 (s, 3H), 1.24 (m, 12H), 1.12 (s, 3H), 1.00 (m, 1H), 0.66 (m, 1H), 0.29 (m, 1H); ¹³C-NMR (CDCl₃) δ 164.45, 153.34, 139.91, 108.17, 71.76, 68.54, 64.23, 47.51, 23.77, 22.45, 17.43, 15.87, 13.87, 12.99.

(±)-trans-1-[2-(Diisopropoxy-phosphorylmethoxymethyl)-cyclopropylmethyl ester]-uracil (15)

Compound **15** was prepared from **11** by following the method as described for **12**: the yield of **15** was 44%: ¹H-NMR (CDCl₃, 300 MHz) δ 7.67 (d, *J* = 8.1 Hz, 1H), 5.66

(d, *J* = 8.2 Hz, 1H), 4.87 (m, 2H), 4.05 (dd, *J* = 15.9, 6.6 Hz, 2H), 3.71 (d, *J* = 8.2 Hz, 2H), 3.66 (d, *J* = 9.9 Hz, 1H), 4.40 (dd, *J* = 15.0, 8.8 Hz, 1H), 3.09 (d, *J* = 11.1 Hz, 1H), 1.32 (m, 12H), 1.10 (s, 3H), 0.98 (m, 1H), 0.52 (m, 1H), 0.23 (m, 1H); ¹³C-NMR (CDCl₃) δ 163.53, 151.12, 143.35, 101.82, 71.02, 70.87, 66.45, 47.67, 23.67, 22.64, 19.17, 15.49, 15.09.

(±)-trans-9-[2-(Methoxymethyl)-2-methyl-cyclopropylmethyl-phosphonic acid]-adenine (16)

To a solution of phosphonate **12** (157 mg, 0.382 mmol) in 10 mL of anhydrous methylene chloride (CH₂)₃SiBr (0.63 g, 4.16 mmol) was added. The mixture was refluxed for overnight and concentrated *in vacuo*. The residue was partitioned between distilled water and washed out by CH₂Cl₂. The aqueous layer was dried by freeze dryer to give **16** (96 mg, 77%) as a solid: UV (H₂O) λ_{max} 261.5 nm; ¹H-NMR (DMSO-*d*₆, 300 MHz) δ 8.02 (s, 1H), 7.70 (s, 1H), 4.07 (dd, *J* = 14.1, 6.6 Hz, 1H), 3.82 (dd, *J* = 13.8, 8.4 Hz, 1H), 3.71 (d, *J* = 8.1 Hz, 2H), 2.78 (m, 2H), 1.11 (s, 3H), 0.97 (m, 1H), 0.57 (m, 1H), 0.21 (m, 1H); ¹³C-NMR (DMSO-*d*₆) δ 154.94, 152.42, 142.93, 115.87, 68.45, 65.82, 49.57, 23.10, 19.72, 15.50, 14.92.

(±)-trans-1-[2-(Methoxymethyl)-2-methyl-cyclopropylmethyl-phosphonic acid]-cytosine (17)

Compound **17** was prepared from **13** by following the method as described for **16**: the yield of **17** was 80%: UV (H₂O) λ_{max} 272.0 nm; ¹H-NMR (DMSO-*d*₆, 300 MHz) δ 7.43 (d, *J* = 6.9 Hz, 1H), 6.69 (br d, 2H), 5.42 (d, *J* = 6.9 Hz, 1H), 3.91 (dd, *J* = 10.8, 5.1 Hz, 1H), 3.51 (dd, *J* = 14.1, 7.2 Hz, 1H), 3.30-2.85 (m, 2H), 1.00-0.83 (m, 4H), 0.55 (dd, *J* = 8.4, 3.9 Hz, 1H), 0.21 (t, *J* = 5.1 Hz, 1H); ¹³C-NMR (DMSO-*d*₆) δ 165.81, 156.02, 145.42, 92.99, 68.69, 64.65, 48.25, 22.29, 19.68, 15.56, 14.67.

(±)-trans-1-[2-(Methoxymethyl)-2-methyl-cyclopropylmethyl phosphonic acid]-thymine (18)

Compound **18** was prepared from **14** by following the method as described for **16**: the yield of **18** was 72%: UV (H₂O) λ_{max} 267.0 nm; ¹H-NMR (DMSO-*d*₆, 300 MHz) δ 7.74 (s, 1H), 3.98 (dd, *J* = 14.0, 8.4 Hz, 1H), 3.71 (d, *J* = 8.2 Hz, 2H), 3.47 (dd, *J* = 14.5, 9.0 Hz, 1H), 3.02 (m, 2H), 1.92 (s, 3H), 1.12-0.99 (m, 4H), 0.63 (m, 1H), 0.30 (t, *J* = 5.6 Hz, 1H); ¹³C-NMR (DMSO-*d*₆) δ 164.67, 154.23, 140.65, 105.23, 68.54, 65.27, 46.35, 22.99, 17.71, 15.12, 14.89, 13.02.

(±)-trans-1-[2-(Methoxymethyl)-2-methyl-cyclopropylmethyl phosphonic acid]-uracil (19)


Compound **19** was prepared from **15** by following the method as described for **16**: the yield of **19** was 74%: UV (H₂O) λ_{max} 262.0 nm; ¹H-NMR (DMSO-*d*₆, 300 MHz) δ

7.45 (d, $J = 7.8$ Hz, 1H), 5.30 (d, $J = 7.7$ Hz, 1H), 3.76 (d, $J = 7.9$ Hz, 2H), 3.55 (dd, $J = 13.8, 81.$ Hz, 1H), 3.34 (dd, $J = 14.4, 8.1$ Hz, 1H), 2.96 (dd, $J = 11.1, 6.0$ Hz, 1H), 2.84 (dd, $J = 11.1, 4.5$ Hz, 1H), 1.02-0.98 (m, 4H), 0.59 (m, 1H), 0.29 (t, $J = 5.2$ Hz, 1H); $^{13}\text{C-NMR}$ (DMSO- d_6) δ 163.70, 151.04, 145.24, 100.67, 68.48, 64.87, 47.38, 22.49, 19.38, 15.52, 14.62.

RESULTS AND DISCUSSION

For the synthesis of target cyclopropyl nucleoside phosphonic acid, acetol **6** was selected as a starting material. As can be seen from Scheme 1, the synthetic route applied for the synthesis of target nucleosides is very simple and straightforward. Allylic alcohol **7** was readily synthesized by following the previously reported procedure (Hong *et al.*, 2003), which was subsequently

subjected to Simmons-Smith carbene cycloaddition (Zhao *et al.*, 1995) in the presence of ZnEt_2 and CH_2I_2 to form cyclopropyl alcohol **8**. The hydroxyl group of **8** was phosphonated by treating it with diisopropyl bromomethylphosphonate in the presence of anhydrous DMF to form the key intermediate **9**. Silyl protection group was readily removed by the treatment with tetrabutylammonium fluoride (TBAF) to form **10**. Compound **10** was methanesulfonylated in the presence of MsCl and TEA in anhydrous CH_2Cl_2 to form **11**, which was coupled with natural bases (adenine, cytosine, thymine, uracil) under well-known classical nucleophilic $\text{S}_{\text{N}}2$ substitution conditions (Hossain *et al.*, 1996) to form the acyclic nucleoside phosphonates **12-15**, respectively. Isopropyl groups of phosphonates were readily hydrolyzed by treatment with trimethylsilylbromide (CH_3SiBr) (El Subbagh *et al.*, 1996) to form final nucleoside phosphonic acids, **16-19**.



Reagents: i) CH_2I_2 , ZnEt_2 , CH_2Cl_2 ; ii) Diisopropyl bromomethylphosphonate, LiOt-Bu , LiI , DMF; iii) TBAF, THF; iv) MsCl , TEA, CH_2Cl_2 ; v) Bases, K_2CO_3 , 18-C-6, DMF; vi) $(\text{CH}_3)_3\text{SiBr}$, CH_2Cl_2 .

Scheme 1. Synthesis of cyclopropyl phosphonic acid nucleosides

Table I. The antiviral activities of the synthesized compounds

Compounds	HIV-1 EC_{50} ($\mu\text{g/mL}$)	HSV-1 EC_{50} ($\mu\text{g/mL}$)	HCMV EC_{50} ($\mu\text{g/mL}$)	CoxB3 EC_{50} ($\mu\text{g/mL}$)	cytotoxicity IC_{50} ($\mu\text{g/mL}$)
16	>100	>100	>100	>100	>100
17	55.7	>100	>100	43.56	>100
18	>100	>100	>100	>100	>100
19	>100	>100	>100	>100	>100
AZT	0.0007	ND	ND	ND	0.5
Ganciclovir	ND	1.24	ND	ND	>10
Ribavirin	ND	ND	ND	30.43	>300

ND: Not Determined

The antiviral assays of the target nucleosides were performed against HIV-1, HSV-1, HCMV, and CoxB3 and the results are shown in Table I. None of the compound exhibited excellent antiviral activity. In the case of cytosine analogue **17**, weak activities against HIV-1 and CoxB3 were observed, without any significant toxicity to the host cell.

To conclude, we have successfully synthesized novel methyl branched cyclopropyl nucleoside phosphonic acids by employing acetol as a starting material and by following Simmons-Smith reaction as a key step. However, when compared with other nucleoside derivatives, cytosine derivative **17** exhibited weak anti-HIV-1 and CoxB3 activity. The results derived from the present study will be useful for the further development of novel cyclopropyl nucleoside phosphonic acids.

REFERENCES

- Ashton, A. T., Meurer, L. C., Cantone, C. L., Field, A. K., Hannah, J., Karkas, J. D., Liou, R., Patel, G. F., Perry, H. C., Wagner, A. F., Walton, E., and Tolman, R. L., Synthesis and antitherapeutic activity of (\pm)-9-[[*Z*]-2-(Hydroxymethyl)cyclopropyl]methyl] guanine and related compounds. *J. Med. Chem.*, 31, 2304-2315 (1988).
- Balzarini, J., Naesens, L., Herdewijn, P., Rosenberg, I., Holy, A., Pauwels, R., Baba, M., Johns, D. G., and De Clercq, E., Marked *in vivo* anti-retrovirus activity of 9-(2-phosphonomethoxyethyl)-adenine, a selective anti-human immunodeficiency virus agent. *Proc. Natl. Acad. Sci. U.S.A.*, 86, 332-336 (1989).
- Bischofberger, N. and Jones, R. J., Nucleotide prodrugs. *Antiviral Res.*, 27, 1-17 (1995).
- De Clercq, E., Sakuma, T., Baba, M., Pauwels, R., Balzarini, J., Rosenberg, I., and Holy, A., Antiviral activity of phosphonyl-methoxyalkyl derivatives of purine and pyrimidines. *Antiviral Res.*, 8, 262-272 (1987).
- El Subbagh, H. I., Racha, S., Abushanab, E., and Panzica, R. P., Synthesis of phosphonate isosteres of 2'-deoxy-1',2'-seco-nucleosides. *J. Org. Chem.*, 61, 890-894 (1996).
- Hong, J. H. and Ko O. H., Synthesis and antiviral evaluation of novel acyclic nucleoside. *Bull. Korean Chem. Soc.*, 24, 1284-1288 (2003).
- Hossain, N., Rozenski, J., De Clercq, E., and Herdewijn, P., Synthesis and antiviral activity of acyclic analogues of 1,5-anhydrohexitol nucleosides using Mitsunobu reaction. *Tetrahedron*, 52, 13655-13670 (1996).
- Noble, S. and Goa, L. L., Adefovir. *Drugs*, 58, 479-487 (1999).
- Qiu, Y. L., Hempel, A., Camerman, N., Camerman, A., Geiser, F., Ptak, R. G., Brietenbach, J. M., Kira, T., Li, L., Gullen, E., Cheng, Y. C., Drach, J. C., and Zemlicka J., (*R*)-(-)- and (*S*)-(+)-synadenol : Synthesis, absolute configuration, and enantioselectivity of antiviral effect. *J. Med. Chem.*, 41, 5257-5264 (1998a).
- Qiu, Y. L., Ksebati, M. B., Ptak, R. G., Fan, B. Y., Breitenbach, J. M., Lin, J. S., Cheng, Y. C., Kern, E. R., Drach, J. C., and Zemlicka, J., (*Z*)- and (*E*)-2-((Hydroxymethyl)-cyclopropylidene) methyladenine and -guanine: New nucleoside analogues with a broad-spectrum antiviral activity. *J. Med. Chem.*, 41, 10-23 (1998b).
- Sekiyama, T., Hatsuya, S., Tanaka, Y., Uchiyama, M., Ono, N., Iwayama, S., Oikawa, M., Suzuki, K., Okunishi, M., and Tsuji, T., Synthesis and antiviral activity of novel acyclic nucleosides: Discovery of a cyclopropyl nucleoside with potent inhibitory activity against herpes viruses. *J. Med. Chem.*, 41, 1284-1298 (1998).
- Snoeck, R., Sakuma, T., De Clercq, E., Rosenberg, I., and Holy, A., (*S*)-1-(3-Hydroxy-2-phosphonyl methoxypropyl)cytosine, a potent and selective inhibitor of human cytomegalovirus replication. *Antimicrob. Agents Chemother.*, 32, 1839-1844 (1988).
- Starrett, J. E., Tortolani, D.R., Russell, J., Hitchcock, M. J. M., Whiterock, V., Martin, J. C., and Mansuri, M. M., Synthesis, oral bioavailability determination and *in vitro* evaluation of prodrugs of the antiviral agent 9-[2-(phosphonomethoxy)ethyl]adenine (PMEA). *J. Med. Chem.*, 37, 1857-1864 (1994).
- Zhao, Y., Yang, T., Lee, M. G., Lee, D. W., Gary Newton, M., and Chu, C. K., Asymmetric synthesis of (1'*S*,2'*R*)-cyclopropyl carbocyclic nucleosides. *J. Org. Chem.*, 60, 5236-5242 (1995).