

NOTES

SYNTHESIS OF ^{14}C -LABELLED TRANEXAMIC ACID[*trans*-AMINO-(^{14}C -METHYL)-CYCLOHEXANE CARBOXYLIC ACID]





Received on June 3, 1975

INTRODUCTION

Tranexamic acid (*trans*-4-aminomethylcyclohexane carboxylic acid) is a valuable antifibrinolytic agent which has been used clinically for several years⁽¹⁾. For biomedical and pharmacological studies of this drug the synthesis of the radioactively labelled compound was required. Although tritium labelling appeared convenient, a metabolically stable ^{14}C -label was preferred due to the nature of the planned investigations, and we now report a procedure for the specific introduction of ^{14}C in the methylene group of the side chain, utilizing potassium ^{14}C -cyanide as the radioactive starting material.

DISCUSSION AND RESULTS

A convenient synthetic route leading to tranexamic acid consists in the reduction of *p*-cyanobenzoic acid⁽²⁾. Introduction of a radioactive *p*-cyano group into benzoic acid thus appeared to be an attractive step in the synthesis of ^{14}C -labelled tranexamic acid. The synthetic route is depicted below:



The conversion of potassium cyanide to cuprous cyanide according to Vogel⁽³⁾ proceeded smoothly also on a small scale. Several solvents have been suggested for the nucleophilic exchange of the *p*-halogen atom of benzoic acids^(4,5,6). In our case the best yield was obtained in dimethylsulphoxide to which a trace of pyridine had been added. *p*-Iodobenzoic acid was found to afford a slightly higher yield of nitrile than the bromo analog (cf.7).

The reduction of *p*-cyanobenzoic acid to 4-aminomethylcyclohexane carboxylic acid in one step has been reported⁽⁸⁾. In our hands, however, better yields and purer products were obtained if the reduction was carried out in two separate steps. The cyano group was first hydrogenated using a Raney-nickel catalyst at atmospheric pressure⁽⁸⁾, whereafter the aromatic ring was reduced in acetic acid over platinum oxide at slightly elevated pressure and temperature⁽²⁾. This procedure afforded predominantly the *cis*-isomer (70-90% and 10-30% *trans*). Isomerization of this mixture in sodium hydroxide solution at 210°⁽⁹⁾ afforded a product with a *trans/cis* ratio of approximately 9 to 1. Crystallization of the amino acid as its toluenesulphonic acid salt from acetone removed the last traces of the *cis*-form. The main fraction of *p*-aminomethylcyclohexane carboxylic acid thus obtained consisted of 98.8% of the *trans*-isomer.

The overall yield based on the radioactive potassium cyanide was about 12%, which appears satisfactory considering the numerous steps involved.

EXPERIMENTAL

Analysis

Separation of the isomers using high voltage paper electrophoresis was carried out at pH 4.5 (50 V, 3 h). The electropherograms were developed with ninhydrine. The ratio of *trans/cis*-isomers and residual *p*-aminomethylbenzoic acid was determined by scanning the strips in a Packard Chromatogram scanner Model 7201. Specific radioactivity was determined by liquid scintillation counting, using a Packard 3375 instrument.

¹⁴C-Copper(I)cyanide

Powdered crystallized copper(I)sulphate (535 mg) was dissolved in water (1.7 ml) in a water-manteled beaker warmed to 60° (the pH should be 3.5 - 4.5). To this solution was added within 1 min sodium hydrogen sulphite (150 mg in 1 ml of water) immediately followed by potassium ¹⁴C-cyanide (150 mg, 100 mCi, in 1 ml of water; The Radiochemical Centre Ltd, Amersham, England) both

solutions warmed to approximately 60°. The resulting mixture was kept at this temperature for 15 min and then filtered through a sintered glass plate. The product was washed with boiling water and by ethanol and finally dried at 110° for 24 h. The yield was 180 mg (87%, based on KCN).

p-(¹⁴C-Cyano)-benzoic acid

p-Iodobenzoic acid (540 mg, 2.2 mM) was dissolved in dimethyl-sulphoxide (10 ml) and to the solution was added the foregoing cuprous-¹⁴C-cyanide (180 mg, 2 mM) followed by 10 drops of pyridine. The mixture was refluxed for 4 h and then poured into a solution of iron(III)chloride (800 mg) in 2.5 N hydrochloric acid (60 ml). The mixture was kept at 60-70° for 20 min in order to decompose the nitrile-copper iodide complex⁽⁵⁾, cooled and extracted with ether (2 x 100 ml). The extracts were dried (Na₂SO₄) and the solvent evaporated. The residue was crystallized from water, yielding 205 mg (70%) of p-(¹⁴C-cyano)-benzoic acid, mp 212-214°, lit.⁽¹⁰⁾ 213-215°.

p-Amino(¹⁴C-methyl)benzoic acid

The p-¹⁴C-cyanobenzoic acid (205 mg) was dissolved in 20% ammonia (10 ml). Raney-nickel (50 mg, obtained from Liljeholmens Stearinfabrik, Stockholm) was added and the mixture was hydrogenated at room temperature and atmospheric pressure until the consumption of hydrogen ceased (~ 2 h). The catalyst was filtered off, the solution was evaporated to dryness and the acid was crystallized from a small volume of water, yielding 140 mg (60%), mp 343°, lit.⁽²⁾ 345°.

4-Amino-(¹⁴C-methyl)cyclohexane carboxylic acid

The labelled 4-aminomethylbenzoic acid (140 mg) was mixed with inactive material (150 mg) and dissolved in acetic acid (10 ml). Five drops of water and 70 mg of PtO₂ were added and the mixture was hydrogenated at 70° and 4 atm for 12 h with 70 mg of PtO₂ as a catalyst. The catalyst was filtered off and the solvent removed, yielding 230 mg (56%) of amino acid acetate (predominantly *cis*-isomer).

Trans-4-amino-(¹⁴C-methyl)-cyclohexanecarboxylic acid. [¹⁴C-Tranexamic acid]

The above *cis-trans* mixture, sodium hydroxide (80 mg) and water (2 ml) were heated at 210° for 20 h in a sealed steel container of 7 ml volume. After cooling, water (10 ml) was added and the resulting solution was passed through a strong cation exchange column (0.5 x 10 cm, Dowex 50 x 8, NH₄⁺). The eluate and washings (5 ml) were evaporated and to the residue was added a solution of p-toluenesulphonic acid (250 mg) in water (20 ml). The resulting mixture was evaporated in vacuo at 50° and the remaining semi-solid was dissolved in acetone (50 ml) whilst still warm. From the slightly turbid so-

lution the *p*-toluenesulphonic acid salt of tranexamic acid crystallized. The solid was collected and dissolved in water (25 ml). This solution was passed through an anion exchange column (1 x 12 cm; Amberlite IR-4B, OH⁻). The eluate was evaporated and the residue was crystallized from water. A first yield of 63 mg (38%) of crystalline *trans*-4-amino-(¹⁴C-methyl)-cyclohexane carboxylic acid was obtained. Electrophoresis showed that this product consisted of 98.8% of the *trans*-isomer. From the mother liquor, a second semi-crystalline fraction was obtained; 30 mg (18%), 94.5% *trans*.

The specific activity of the ¹⁴C-tranexamic acid was 20.4 mCi/mM. The total recovery of radioactivity in the two fractions was 12.4 mCi (= 12.4% overall yield based on K¹⁴CN).

ACKNOWLEDGEMENT

The authors wish to thank Mr. O.Eriksson and Mr. J. Friberg, Research Department, AB Kabi, for performing the analysis.

Jan Lundström^x

Gertrud Westin-Sjödahl^{xx} and N. Åke Jönsson

^xFaculty of Pharmacy, Biomedical Center, University of Uppsala

Box 570, S-751 23 Uppsala. Present address: Astra Läkemedel,
S-151 85 Södertälje, Sweden

^{xx}Department of Organic Chemistry, Research Department, AB Kabi, Fack, S-104 25
Stockholm, Sweden. To whom requests for reprints should be sent.

REFERENCES

1. Andersson L. - S.A.Med.J. 46:182 (1972)
2. Levine M. and Sedlecky R. - J.Org.Chem.24:115 (1959)
3. Vogel A.J. - "Practical Organic Chemistry", 3rd edn, p 191, Longmans, London (1961)
4. Friedman L. and Shechter H. - J.Org.Chem.26:2522 (1961)
5. Newman M.S. and Boden H. - J.Org.Chem.26:2525 (1961)
6. Bacon R.G.R. and Hill H.A.O. - Proc.Chem.Soc.1962:113
7. Bacon R.G.R. and Hill H.A.O. - J.Chem.Soc.1964:1097
8. Albert A. and Magrath D. - J.Chem.Soc.1944:678
9. Naito T., Okamo J., Inaoka M., Kadoya S., Shindo M. and Aoyazi T. - Jap.pat. 14,830, cf C.A.65:p 200326 (1966)
10. Handbook of Chemistry and Physics, 41st edn. Chemical Rubber Co. (1959-60)